



533 Rec'd PCT/PTO 21 AUG 2001

09/913980

TRANSMITTAL LETTER TO THE UNITED STATES

ATTORNEY'S DOCKET NUMBER 49774

DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

INTERNATIONAL APPLICATION NO.  
PCT/EP00/01070

INTERNATIONAL FILING DATE  
10 February 2000

PRIORITY DATE CLAIMED  
22 February 1999

TITLE OF INVENTION: HAIR COSMETIC FORMULATIONS

APPLICANT(S) FOR DO/EO/US Reinhold DIEING, Michael GOTSCHKE, Peter HOESSEL, Axel SANNER,  
Alfred LEINENBACH, Keith Leslie RUTHERFORD

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. /X/ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
  2. / / This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
  3. /X/ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
  4. /x / A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
  5. /X/ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
    - a. /X/ is transmitted herewith (required only if not transmitted by the International Bureau).
    - b. / / has been transmitted by the International Bureau.
    - c. / / is not required, as the application was filed in the United States Receiving Office (RO/USO).
  6. /X/ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
  7. / / Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
    - a. / / are transmitted herewith (required only if not transmitted by the International Bureau).
    - b. / / have been transmitted by the International Bureau.
    - c. / / have not been made; however, the time limit for making such amendments has NOT expired.
    - d. / / have not been made and will not be made.
  8. / / A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
  9. / X / An oath or declaration of the inventor(s) (35 U.S.C. 171(c)(4)).
  10. / / A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern other document(s) or information included:
11. / X / An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
  12. /x / An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
  13. /X / A FIRST preliminary amendment.  
/ / A SECOND or SUBSEQUENT preliminary amendment.
  14. / / A substitute specification.
  15. / / A change of power of attorney and/or address letter.
  16. /x / Other items or information.  
International Search Report  
International Preliminary Examination Report

09/913980

U.S. Appl. No. (If Known) INTERNATIONAL APPLN. NO.  
PCT/EP00/01070

ATTORNEY'S DOCKET NO.  
49774

	CALCULATIONS	PTO USE ONLY
17. /X/ The following fees are submitted		
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):		
Search Report has been prepared by the		
EPO or JPO.....\$860.00	860.00	
International preliminary examination fee paid to USPTO		
(37 CFR 1.482).....\$750.00		
No international preliminary examination fee paid to		
USPTO (37 CFR 1.482) but international search fee paid		
to USPTO (37 CFR 1.445(a)(2)).....\$700.00		
Neither international preliminary examination fee		
(37 CFR 1.482) nor international search fee		
(37 CFR 1.445(a)(2)) paid to USPTO .....\$ 970.00		
International preliminary examination fee paid to		
USPTO (37 CFR 1.482) and all claims satisfied pro		
-visions of PCT Article 33(2)-(4).....\$96.00		

ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 860.00

Surcharge of \$130.00 for furnishing the oath or declaration  
later than / / 20 / / 30 months from the earliest  
claimed priority date (37 CFR 1.492(e)).

Claims	Number Filed	Number Extra	Rate	
Total Claims	31 -20	11	X\$18.	198.00
Indep.Claims	5 -3	2	X\$80.	160.00
Multiple dependent claim(s)(if applicable)		+270.		
TOTAL OF ABOVE CALCULATION			=	1,218.00

Reduction of 1/2 for filing by small entity, if applicable.  
Verified Small Entity statement must also be filed  
(Note 37 CFR 1.9, 1.27, 1.28).

SUBTOTAL = 1,218.00

Processing fee of \$130. for furnishing the English  
translation later than / / 20 / / 30 months from the  
earliest claimed priority date (37 CFR 1.492(f)).

TOTAL NATIONAL FEE = 1,218.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)).  
The assignment must be accompanied by an appropriate cover  
sheet (37 CFR 3.28, 3.31) \$40.00 per property =

TOTAL FEES ENCLOSED = \$ 1,258.00

Amount to be  
refunded: \$  
Charged \$

- a./X/ A check in the amount of \$ 1,258. to cover the above fees is enclosed.
- b./ / Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c./X/ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 11-0345. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:  
KEIL & WEINKAUF  
1101 Connecticut Ave., N.W.  
Washington, D. C. 20036

SIGNATURE

Herbert B. Keil

NAME

Registration No. 18,967

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of )  
DIEING et al. ) BOX-PCT  
)  
International Application )  
PCT/EP 00/01070 )  
)  
Filed: February 10, 2000 )  
)

For: HAIR COSMETIC FORMULATIONS

PRELIMINARY AMENDMENT

Honorable Commissioner of  
Patents and Trademarks  
Washington, D.C. 20231

Sir:

Prior to examination, kindly amend the above-identified application as follows:

IN THE CLAIMS

Amend the claims as shown in the attached sheets.

R E M A R K S

The claims have been amended to eliminate multiple dependency and to place them in better form for U.S. filing. No new matter is included.

A clean copy of the claims is attached.

Favorable action is solicited.

Respectfully submitted,

KEIL & WEINKAUF



Herbert B. Keil  
Reg. No. 18,967

1101 Connecticut Ave., N.W.  
Washington, D.C. 20036

(202)659-0100

09/913980

1900	1901	1902	1903	1904	1905	1906	1907	1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929	1930	1931	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

10. The use of polymers as claimed in claim 8, wherein the alkylene oxide used is ethylene oxide.
11. The use of polymers as claimed in claim 8, wherein the polyethyleneimine has a molecular weight between 300 and 20000.
15. The use of polymers as claimed in claim 1, wherein c) is chosen from the group: acrylic acid, methacrylic acid, maleic acid, fumaric acid, crotonic acid, maleic anhydride and its half-esters, methyl acrylate, methyl methacrylate, ethyl acrylate, ethyl methacrylate, n-butyl acrylate, n-butyl methacrylate, t-butyl acrylate, t-butyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, stearyl acrylate, stearyl methacrylate, N-t-butylacrylamide, N-octylacrylamide, 2-hydroxyethyl acrylate, hydroxypropyl acrylates, 2-hydroxyethyl methacrylate, hydroxypropyl methacrylates, alkylene glycol (meth)acrylates, styrene, unsaturated sulfonic acids such as, for example, acrylamidopropane sulfonic acid, vinyl pyrrolidone, vinyl caprolactam, vinyl ethers, (e.g. methyl, ethyl, butyl or dodecyl vinyl ethers), vinylformamide, vinylmethacetamide, vinylamine, 1-vinylimidazole, 1-vinyl-2-methylimidazole, N,N-dimethylaminomethyl methacrylate and N-[3-(dimethylamino)propyl]methacrylamide; 3-methyl-1-vinylimidazolium chloride, 3-methyl-1-vinylimidazolium methylsulfate, N,N-dimethylaminoethyl methacrylate, N-[3-(dimethylamino)propyl]methacrylamide quaternized with methyl chloride, methyl sulfate or diethyl sulfate.

## CLEAN VERSION OF AMENDED CLAIMS - OZ 49774

16. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 10 - 90 % by weight
  - b) 2 - 90 % by weight
  - c) 0 - 50 % by weight.
17. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 50 - 97 % by weight
  - b) 3 - 50 % by weight
  - c) 0 - 30 % by weight.
18. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 60 - 97 % by weight
  - b) 3 - 40 % by weight
  - c) 0 - 20 % by weight.
19. The use as claimed in claim 1, where a crosslinking is carried out after the hydrolysis.

[illegible]

10. The use of polymers as claimed in claim 8 [claims 8 and 9], wherein the alkylene oxide used is ethylene oxide.
11. The use of polymers as claimed in claim 8 [claims 8, 9 and 10], wherein the polyethyleneimine has a molecular weight between 300 and 20000.
15. The use of polymers as claimed in claim 1 [claims 1 to 14], wherein c) is chosen from the group:  
  
acrylic acid, methacrylic acid, maleic acid, fumaric acid, crotonic acid, maleic anhydride and its half-esters, methyl acrylate, methyl methacrylate, ethyl acrylate, ethyl methacrylate, n-butyl acrylate, n-butyl methacrylate, t-butyl acrylate, t-butyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, stearyl acrylate, stearyl methacrylate, N-t-butylacrylamide, N-octylacrylamide, 2-hydroxyethyl acrylate, hydroxypropyl acrylates, 2-hydroxyethyl methacrylate, hydroxypropyl methacrylates, alkylene glycol (meth)acrylates, styrene, unsaturated sulfonic acids such as, for example, acrylamidopropane sulfonic acid, vinyl pyrrolidone, vinyl caprolactam, vinyl ethers, (e.g. methyl, ethyl, butyl or dodecyl vinyl ethers), vinylformamide, vinylmethacetamide, vinylamine, 1-vinylimidazole, 1-vinyl-2-methylimidazole, N,N-dimethylaminomethyl methacrylate and N-[3-(dimethylamino)propyl]methacrylamide;  
  
3-methyl-1-vinylimidazolium chloride, 3-methyl-1-vinylimidazolium methylsulfate, N,N-dimethylaminoethyl methacrylate,  
  
N-[3-(dimethylamino)propyl]methacrylamide quaternized with methyl chloride,


MARKED UP VERSION OF AMENDED CLAIMS - OZ 49774

methyl sulfate or diethyl sulfate.

16. The use of polymers as claimed in claim 1 [claims 1 and 15], wherein the quantitative ratios are
- a) 10 - 90 % by weight
  - b) 2 - 90 % by weight
  - c) 0 - 50 % by weight.
17. The use of polymers as claimed in claim 1 [claims 1 to 15], wherein the quantitative ratios are
- a) 50 - 97 % by weight
  - b) 3 - 50 % by weight
  - c) 0 - 30 % by weight.
18. The use of polymers as claimed in claim 1 [claims 1 to 15], wherein the quantitative ratios are
- a) 60 - 97 % by weight
  - b) 3 - 40 % by weight
  - c) 0 - 20 % by weight.
19. The use as claimed in claim 1 [claims 1 to 18], where a crosslinking is carried out after the hydrolysis.

# CURRENT CLAIMS - OZ 49774

1. The use of polymers obtainable by free-radical polymerization of
  - a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of
  - b) polyether-containing compounds and
  - c) optionally one or more other copolymerizable monomers
 and subsequent at least partial hydrolysis of the ester functions of the original monomers a), in hair cosmetic formulations.
2. The use of polymers as claimed in claim 1, wherein the polymers are obtainable by free-radical polymerization of
  - a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of
  - b) polyether-containing compounds of the formula I



in which the variables independently of one another have the following meanings:

R<sup>1</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-, polyalcohol radical;

R<sup>5</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;



## CURRENT CLAIMS - OZ 49774

R<sup>2</sup> to R<sup>4</sup> are

-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -CH<sub>2</sub>-CH(R<sup>6</sup>)-, -CH<sub>2</sub>-CHOR<sup>7</sup>-CH<sub>2</sub>-;

R<sup>6</sup> is C<sub>1</sub>-C<sub>24</sub>-alkyl;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

A is -C(=O)-O-, -C(=O)-B-C(=O)-O-,

-C(=O)-NH-B-NH-C(=O)-O-;

B is -(CH<sub>2</sub>)<sub>t</sub>-, arylene, optionally substituted;

n is from 1 to 1000;

s is from 0 to 1000;

t is from 1 to 12;

u is from 1 to 5000;

v is from 0 to 5000;

w is from 0 to 5000;

x is from 0 to 5000;

y is from 0 to 5000;

z is from 0 to 5000;

and

c) optionally one or more other copolymerizable monomers

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

3. The use of polymers as claimed in claim 2, wherein the polymers are obtainable

## CURRENT CLAIMS - OZ 49774

by free-radical polymerization of

- a) at least one vinyl ester of  $C_1$ - $C_{24}$ -carboxylic acids in the presence of
- b) polyether-containing compounds of the formula I having an average molecular weight of from 300 to 100000 (number average), in which the variables independently of one another have the following meanings:

$R^1$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-, polyalcohol radical;

$R^5$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

$R^2$  to  $R^4$  are

$-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH_2-CH(R^6)-$ ,  $-CH_2-CHOR^7-CH_2-$ ;

$R^6$  is  $C_1$ - $C_{12}$ -alkyl;

$R^7$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

n is from 1 to 8;

s is 0;

u is from 2 to 2000;

v is from 0 to 2000;

w is from 0 to 2000;

and

- c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

## CURRENT CLAIMS - OZ 49774

4. The use of polymers as claimed in claim 2, wherein the polymers are obtainable by free-radical polymerizable of

- a) at least one vinyl ester of  $C_1$ - $C_{24}$ -carboxylic acids in the presence of
- b) polyether-containing compounds of the formula I having an average molecular weight of from 500 to 50000 (number average), in which the variables independently of one another have the following meaning:

$R^1$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

$R^5$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

$R^2$  to  $R^4$  are

$-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH_2-CH(R^6)-$ ,  $-CH_2-CHOR^7-CH_2-$ ;

$R^6$  is  $C_1$ - $C_6$ -alkyl;

$R^7$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

n is 1;

s is 0;

u is from 5 to 500;

v is from 0 to 500;


w is from 0 to 500;

and


- c) optionally at least one or more other copolymerizable monomers and subsequent at least partial hydrolysis of the ester functions of the original monomers a), in hair cosmetic formulations.

# CURRENT CLAIMS - OZ 49774





5. The use of polymers as claimed in claim 1, wherein the polymers are obtainable by free-radical polymerization of
  - a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of
  - b) polyether-containing silicone derivatives
  - and
  - c) optionally one or more other copolymerizable monomers
 and subsequent at least partial hydrolysis of the ester function of the original monomers a).
6. The use of polymers as claimed in claim 5, wherein the polymers are obtainable by free-radical polymerization of
  - a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of
  - b) polyether-containing silicone derivatives of the formula II




where:



# CURRENT CLAIMS - OZ 49774



$R^{13}$  is a  $C_1$ - $C_{40}$  organic radical which can contain amino, carboxyl or sulfonate groups, or where  $e = 0$ , is also the anion of an inorganic acid, and where the radicals  $R^8$  can be identical or different, and come either from the group of aliphatic hydrocarbons having from 1 to 20 carbon atoms, are cyclic aliphatic hydrocarbons having from 3 to 20 carbon atoms, are of an aromatic nature or are identical to  $R^{12}$ , where:



# CURRENT CLAIMS - OZ 49774

with the proviso that at least one of the radicals  $R^8$ ,  $R^9$  or  $R^{10}$  is a polyalkylene oxide-containing radical as defined above,

and  $f$  is an integer from 1 to 6,


$a$  and  $b$  are integers such that the molecular weight of the polysiloxane block is between 300 and 30000,

$c$  and  $d$  can be integers between 0 and 50, with the proviso that the sum  $c + d$  is greater than 0, and  $e$  is 0 or 1,

and

optionally one or more other copolymerizable monomers and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

7. The use of polymers as claimed in claim 6, wherein formula II has the following meaning:



8. The use as claimed in claim 1, wherein the polymers are obtainable by free-radical polymerization of

## CURRENT CLAIMS - OZ 49774

- a) at least one vinyl ester of  $C_1$ - $C_{24}$ -carboxylic acids in the presence of
  - b) polyether-containing compounds obtainable by reaction of  
polyethyleneimines with alkylene oxides
  - and
  - c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).
9. The use of polymers as claimed in claim 8, wherein the alkylene oxides used are ethylene oxide, propylene oxide, butylene oxide and mixtures thereof.
  10. The use of polymers as claimed in claim 8, wherein the alkylene oxide used is ethylene oxide.
  11. The use of polymers as claimed in claim 8, wherein the polyethyleneimine has a molecular weight between 300 and 20000.
  12. The use of polymers as claimed in claim 1, wherein the polyether-containing compounds b) have been prepared by polymerization of ethylenically unsaturated alkylene oxide-containing monomers and optionally other copolymerizable monomers.
  13. The use of polymers as claimed in claim 12, wherein the polyether-containing compounds b) have been prepared by polymerization of polyalkylene oxide vinyl ethers and optionally other copolymerizable monomers.

## CURRENT CLAIMS - OZ 49774

14. The use of polymers as claimed in claim 12, wherein the polyether-containing compounds b) have been prepared by polymerization of polyalkylene oxide (meth)acrylates and optionally other copolymerizable monomers.
15. The use of polymers as claimed in claim 1, wherein c) is chosen from the group:  
acrylic acid, methacrylic acid, maleic acid, fumaric acid, crotonic acid, maleic anhydride and its half-esters, methyl acrylate, methyl methacrylate, ethyl acrylate, ethyl methacrylate, n-butyl acrylate, n-butyl methacrylate, t-butyl acrylate, t-butyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, stearyl acrylate, stearyl methacrylate, N-t-butylacrylamide, N-octylacrylamide, 2-hydroxyethyl acrylate, hydroxypropyl acrylates, 2-hydroxyethyl methacrylate, hydroxypropyl methacrylates, alkylene glycol (meth)acrylates, styrene, unsaturated sulfonic acids such as, for example, acrylamidopropane sulfonic acid, vinyl pyrrolidone, vinyl caprolactam, vinyl ethers, (e.g. methyl, ethyl, butyl or dodecyl vinyl ethers), vinylformamide, vinylmethylacetamide, vinylamine, 1-vinylimidazole, 1-vinyl-2-methylimidazole, N,N-dimethylaminomethyl methacrylate and N-[3-(dimethylamino)propyl]methacrylamide;  
3-methyl-1-vinylimidazolium chloride, 3-methyl-1-vinylimidazolium methylsulfate, N,N-dimethylaminoethyl methacrylate,  
N-[3-(dimethylamino)propyl]methacrylamide quaternized with methyl chloride, methyl sulfate or diethyl sulfate.



## CURRENT CLAIMS - OZ 49774

16. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 10 - 90 % by weight
  - b) 2 - 90 % by weight
  - c) 0 - 50 % by weight.
17. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 50 - 97 % by weight
  - b) 3 - 50 % by weight
  - c) 0 - 30 % by weight.
18. The use of polymers as claimed in claim 1, wherein the quantitative ratios are
  - a) 60 - 97 % by weight
  - b) 3 - 40 % by weight
  - c) 0 - 20 % by weight.
19. The use as claimed in claim 1, where a crosslinking is carried out after the hydrolysis.
20. The use as claimed in claim 19, where the crosslinking is carried out by aldehydes, dialdehydes or borates.
21. A hair cosmetic formulation which has the following composition:
  - a) 0.05 - 20 % by weight of the polymer as in claim 1
  - b) 20 - 99.95 % by weight of water and/or alcohol
  - c) 0 - 79.05 % by weight of other constituents.

## CURRENT CLAIMS - OZ 49774

22. A hair cosmetic formulation which has the following composition:

- a) 0.1 - 10 % by weight of the polymer as in claim 1
- b) 20 - 99.9 % by weight of water and/or alcohol
- c) 0 - 70 % by weight of a propellant
- d) 0 - 20 % by weight of other constituents.

23. A hair cosmetic formulation which has the following composition:

- a) 0.1 - 10 % by weight of the polymer as in claim 1
- b) 55 - 94.8 % by weight of water and/or alcohol
- c) 5 - 20 % by weight of a propellant
- d) 0.1 - 5 % by weight of an emulsifier
- e) 0 - 10 % by weight of other constituents.

24. A hair cosmetic formulation which has the following composition:

- a) 0.1 - 10 % by weight of the polymer as in claim 1
- b) 60 - 99.85 % by weight of water and/or alcohol
- c) 0.05 - 10 % by weight of a gel former
- d) 0 - 20 % by weight of other constituents.

25. A hair cosmetic formulation which has the following composition:

- a) 0.05 - 10 % by weight of the polymer as in claim 1,
- b) 25 - 94.95 % by weight of water
- c) 5 - 50 % by weight of surfactants
- d) 0 - 5 % by weight of another conditioning agent

CURRENT CLAIMS - OZ 49774

- e) 0 - 10 % by weight of other cosmetic constituents.

26. A polymer obtainable by free-radical polymerization of
- a) at least one vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid, in the presence of
  - b) polyether-containing silicone derivatives and
  - c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).
27. A polymer obtainable by free-radical polymerization of
- a) a vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid in the presence of
  - b) polyether-containing compounds obtainable by reaction of polyethyleneimines with alkylene oxides and
  - c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).
28. A polymer obtainable by free-radical polymerization of
- a) a vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid in the presence of
  - b) homo- and copolymers of ethylenically unsaturated polyether-containing compounds and
  - c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

CURRENT CLAIMS - OZ 49774

29. A crosslinked polymer obtainable by free-radical polymerization of
- a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub> carboxylic acids in the presence of
  - b) polyether-containing compounds and
  - c) optionally one or more further copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a), where the crosslinker used is either already present during the polymerization, or is added after the polymerization and hydrolysis.
30. A crosslinked polymer as claimed in claim 29, where the crosslinkers used are aldehydes, dialdehydes or borates.
31. The crosslinked polymer as claimed in claim 29, wherein the crosslinker is already present during the polymerization.

52103500

10

It is an object of the present invention to find hair cosmetic  
30 formulations containing film-forming polymers which impart strong  
hold and also high elasticity to the hairstyle.

45

DE 1 077 430 describes a process for the preparation of graft polymers of vinyl esters on polyalkylene glycols.


DE 1 094 457 and DE 1 081 229 describe processes for the preparation of graft polymers of polyvinyl alcohol on polyalkylene glycols by hydrolysis of the vinyl esters and their use as protective colloids, water-soluble packaging films, as sizes and finishes for textiles and in cosmetics.

10 In the preparation of the polymers used according to the invention, grafting onto the polyether-containing compounds (b) can result during the polymerization, which can lead to the advantageous properties of the polymers. However, mechanisms other than grafting are also conceivable.

15 Depending on the degree of grafting, the polymers used according to the invention are taken to mean pure graft polymers and also mixtures of the abovementioned graft polymers with nongrafted polyether-containing compounds and homo- or copolymers of the monomers a) and c).

Polyether-containing compounds (b) which can be used are either polyalkylene oxides, based on ethylene oxide, propylene oxide, butylene oxide and other alkylene oxides, or polyglycerol.

25 Depending on the type of monomer units, the polymers contain the following structural units.




30 where

$\text{R}^6$  is  $\text{C}_1$ - $\text{C}_{24}$ -alkyl;

35  $\text{R}^7$  is hydrogen,  $\text{C}_1$ - $\text{C}_{24}$ -alkyl,  $\text{R}^6-\text{C}(=\text{O})-$ ,  $\text{R}^6-\text{NH}-\text{C}(=\text{O})-$ .

The structural units can either be homopolymers or random copolymers and block copolymers.

40 As polyether (b), preference is given to using polymers of the formula I,



45

I

## 3

in which the variables independently of one another have the following meanings:

5  $R^1$  is hydrogen,  $C_1$ - $C_{24}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-, polyalcohol radicals;

$R^5$  is hydrogen,  $C_1$ - $C_{24}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

10  $R^2$  to  $R^4$  are  
-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -CH<sub>2</sub>-CH( $R^6$ )-, -CH<sub>2</sub>-CHOR<sup>7</sup>-CH<sub>2</sub>-;

$R^6$  is  $C_1$ - $C_{24}$ -alkyl;

15  $R^7$  is hydrogen,  $C_1$ - $C_{24}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

A is -C(=O)-O, -C(=O)-B-C(=O)-O,  
-C(=O)-NH-B-NH-C(=O)-O;

20 B is -(CH<sub>2</sub>)<sub>t</sub>-, arylene, optionally substituted;

n is from 1 to 1000;

s is from 0 to 1000;

25 t is from 1 to 12;

u is from 1 to 5000;

v is from 0 to 5000;

30 w is from 0 to 5000;

x is from 0 to 5000;

35 y is from 0 to 5000;

z is from 0 to 5000.

40 The terminal primary hydroxyl groups of the polyethers prepared on the basis of polyalkylene oxides, and the secondary OH-groups of polyglycerol can in this connection either be present in free unprotected form, or be etherified with alcohols of chain length  $C_1$ - $C_{24}$  or esterified with carboxylic acids of chain length  $C_1$ - $C_{24}$ , or reacted with isocyanates to give urethanes.

45

- Alkyl radicals which may be mentioned for  $R^1$  and  $R^5$  to  $R^7$  are branched or unbranched  $C_1$ - $C_{24}$ -alkyl chains, preferably methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, 2-ethylhexyl, n-octyl, n-nonyl, n-decyl, n-undecyl, n-dodecyl, n-tridecyl, n-tetradecyl, n-pentadecyl, n-hexadecyl, n-heptadecyl, n-octadecyl, n-nonadecyl or n-eicosyl.

Preferred representatives of the abovementioned alkyl radicals which may be mentioned are branched or unbranched  $C_1$ - $C_{12}$ -, particularly preferably  $C_1$ - $C_6$ -alkyl chains.

- The molecular weight of the polyethers is less than 1000000 (number average), preferably in the range from 300 to 100000, particularly preferably in the range from 500 to 50000, very particularly preferably in the range from 800 to 40000.
- Homopolymers of ethylene oxide or copolymers carrying an ethylene oxide content of from 40 to 99% by weight are advantageously used. For the ethylene oxide polymers to be used in preference, the content of copolymerized ethylene oxide is thus from 40 to 100 mol%. Suitable comonomers for these copolymers are propylene oxide, butylene oxide and/or isobutylene oxide. Suitable examples are copolymers of ethylene oxide and propylene oxide, copolymers of ethylene oxide and butylene oxide, and copolymers of ethylene oxide, propylene oxide and at least one butylene oxide. The ethylene oxide content of the copolymers is preferably from 40 to 99 mol%, the propylene oxide content is from 1 to 60 mol% and the content of butylene oxide in the copolymers is from 1 to 30 mol%. As well as straight-chain homo- or copolymers, it is also possible to use branched homo- or copolymers as polyether-containing compounds b).

- Branched polymers can be prepared by adding ethylene oxide and optionally also propylene oxide and/or butylene oxides to, for example, polyalcohol radicals, e.g. to pentaerythritol, glycerol, or to sugar alcohols such as D-sorbitol and D-mannitol, but also to polysaccharides such as cellulose and starch. The alkylene



oxide units can be randomly distributed or be in the form of blocks within the polymer.

It is, however, also possible to use polyesters of polyalkylene  
 5 oxides and aliphatic or aromatic dicarboxylic acids, e.g. oxalic acid, succinic acid, adipic acid and terephthalic acid having molar masses of from 1500 to 25000, as described, for example, in EP-A-0 743 962, as polyether-containing compound. In addition, it is also possible to use polycarbonates by reaction of  
 10 polyalkylene oxides with phosgene or carbonates such as, for example, diphenyl carbonate, and polyurethanes by reaction of polyalkylene oxides with aliphatic and aromatic diisocyanates.

Particularly preferred polyethers (b) are polymers of the formula  
 15 I having an average molecular weight of from 300 to 100000 (number average), in which the variables independently of one another have the following meanings:

$R^1$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-,  
 20 polyalcohol radical;

$R^5$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

$R^2$  to  $R^4$  are  
 25  $-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH_2-CH(R^6)-$ ,  $-CH_2-CHOR^7-CH_2-$ ;

$R^6$  is  $C_1$ - $C_{12}$ -alkyl;

$R^7$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;  
 30

$n$  is from 1 to 8;

$s$  is 0;

35  $u$  is from 2 to 2000;

$v$  is from 0 to 2000;

$w$  is from 0 to 2000.

40 Very particularly preferred polyethers b) are polymers of the formula I having an average molecular weight of from 500 to 50000 (number average), in which the variables independently of one another have the following meanings:

45  $R^1$  is hydrogen,  $C_1$ - $C_6$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

6

R<sup>5</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

R<sup>2</sup> to R<sup>4</sup> are

-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -CH<sub>2</sub>-CH(R<sup>6</sup>)-, -CH<sub>2</sub>-CHOR<sup>7</sup>-CH<sub>2</sub>-;

5

R<sup>6</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

10 n is 1;

s is 0;

u is from 5 to 500;

15


v is from 0 to 500;

w is from 0 to 500.

- 20 However, the polyethers used may also be silicone derivatives. Suitable silicone derivatives are the compounds known under the INCI name dimethicone copolyols or silicone surfactants, such as, for example, those available under the tradenames Abil® (T. Goldschmidt), Alkasil® (Rhône-Poulenc), Silicone Polyol Copolymer® (Genesee), Belsil® (Wacker), Silwet® (Witco, Greenwich, CT, USA) or Dow Corning (Dow Corning). These include compounds having the CAS numbers 64365-23-7; 68937-54-2; 68938-54-5; 68937-55-3.

- 30 Silicones are generally used in hair cosmetics to improve the feel. The use of polyether-containing silicone derivatives as polyether (b) in the polymers according to the invention can therefore additionally lead to an improvement in the feel of the hair.


- 35 Preferred representatives of those polyether-containing silicone derivatives are those which contain the following structural elements:




where:

45

7


 $R^9 = \text{CH}_3$  or

5

 $R^{10} = \text{CH}_3$  or  $R^9$  $R^{11} = \text{H}, \text{CH}_3,$ 

10


15



20  $R^{13}$  is a  $\text{C}_1$ - $\text{C}_{40}$  organic radical which can contain amino, carboxyl or sulfonate groups, or where  $e = 0$ , is also the anion of an inorganic acid,

and where the radicals  $R^8$  can be identical or different, and come  
 25 either from the group of aliphatic hydrocarbons having from 1 to 20 carbon atoms, are cyclic aliphatic hydrocarbons having from 3 to 20 carbon atoms, are of an aromatic nature or are identical to  $R^{12}$ , where:

30



35

with the proviso that at least one of the radicals  $R^8$ ,  $R^9$  or  $R^{10}$  is a polyalkylene oxide-containing radical as defined above, and  $f$  is an integer from 1 to 6,

$a$  and  $b$  are integers such that the molecular weight of the

40 polysiloxane block is between 300 and 30000,

$c$  and  $d$  can be integers between 0 and 50, with the proviso that the sum  $c + d$  is greater than 0, and  $e$  is 0 or 1.

Preferred radicals  $R^9$  and  $R^{12}$  are those in which the sum  $c+d$  is  
 45 between 5 and 30.


## 8

The groups  $R^8$  are preferably chosen from the following group: methyl, ethyl, propyl, butyl, isobutyl, pentyl, isopentyl, hexyl, octyl, decyl, dodecyl and octadecyl, cycloaliphatic radicals, specifically cyclohexyl, aromatic groups, specifically phenyl or naphthyl, mixed aromatic-aliphatic radicals such as benzyl or phenyl ethyl and tolyl and xylyl and  $R^{12}$ .

Particularly suitable radicals  $R^{11}$  are those in which where  $R^{11} = -(\text{CO})_e-R^{13}$   $R^{13}$  is any desired alkyl, cycloalkyl or aryl radical which has between 1 and 40 carbon atoms and which can carry other ionogenic groups such as  $\text{NH}_2$ ,  $\text{COOH}$ ,  $\text{SO}_3\text{H}$ .

Preferred inorganic radicals  $R^{13}$  are, where  $e = 0$ , phosphate and sulfate.

Particularly preferred polyether-containing silicone derivatives are those of the structure:



- In addition, homo- and copolymers of polyalkylene oxide-containing ethylenically unsaturated monomers, such as, for example, polyalkylene oxide (meth)acrylates, polyalkylene oxide vinyl ethers, polyalkylene oxide (meth)acrylamides, polyalkylene oxide allylamides or polyalkylene oxide vinylamides can also be used as polyether (b). It is of course also possible to use copolymers of such monomers with other ethylenically unsaturated monomers.
- It is, however, also possible to use reaction products of polyethyleneimines with alkylene oxides as polyether-containing compounds b). In this case, the alkylene oxides used are preferably ethylene oxide, propylene oxide, butylene oxide and mixtures thereof, particularly preferably ethylene oxide.
- Polyethyleneimines which can be used are polymers having number-average molecular weights of from 300 to 20000, preferably from 500 to 10000, very particularly preferably from 500 to 5000. The weight ratio between used alkylene oxide and polyethyleneimine is in the range from 100 : 1 to 0.1 : 1, preferably in the range from 50 : 1 to 0.5 : 1, very particularly preferably in the range from 20 : 1 to 0.5 : 1.

For the polymerization in the presence of the polyethers b), the following free-radically polymerizable monomers may be mentioned as component a):

- 5 Vinyl esters of aliphatic, saturated or unsaturated  $C_1-C_{24}$  carboxylic acids, such as, for example, formic acid, acetic acid, propionic acid, butyric acid, valeric acid, isovaleric acid, caproic acid, caprylic acid, capric acid, undecylenic acid, lauric acid, myristic acid, palmitic acid, palmitoleic acid,  
 10 stearic acid, oleic acid, arachidic acid, behenic acid, lignoceric acid, cerotinic acid and melissic acid.

Preference is given to using vinyl esters of the abovementioned  $C_1-C_{12}$  carboxylic acids, in particular of  $C_1-C_6$  carboxylic acids.

- 15 Vinyl acetate is very particularly preferred.

It is, of course, also possible to copolymerize mixtures of the respective monomers from group a).

- 20 The vinyl esters (a) can in addition also be used in admixture with one or more ethylenically unsaturated copolymerizable comonomers (c), where the content of these additional monomers should be limited to a maximum of 50% by weight. Preference is given to contents of from 0 to 20% by weight. The term  
 25 ethylenically unsaturated means that the monomers have at least one free-radically polymerizable carbon-carbon double bond which can be mono-, di-, tri- or tetrasubstituted.

The preferred ethylenically unsaturated comonomers (c)

- 30 additionally used can be described by the following formula:



where

35

X is chosen from the group of radicals  $-OH$ ,  $-OM$ ,  $-OR^{16}$ ,  $NH_2$ ,  $-NHR^{16}$ ,  $N(R^{16})_2$ ;

M is a cation chosen from the group consisting of:  $Na^+$ ,  $K^+$ ,  $Mg^{++}$ ,

- 40  $Ca^{++}$ ,  $Zn^{++}$ ,  $NH_4^+$ , alkyl ammonium, dialkylammonium, trialkylammonium and tetraalkylammonium.

The radicals  $R^{16}$  can be identical or different and chosen from the group consisting of  $-H$ ,  $C_1-C_{40}$  linear or branched alkyl radicals,

- 45  $N,N$ -dimethylaminoethyl, 2-hydroxyethyl, 2-methoxyethyl, 2-ethoxyethyl, hydroxypropyl, methoxypropyl or ethoxypropyl.

R<sup>15</sup> and R<sup>14</sup> are independently of one another chosen from the group consisting of: -H, C<sub>1</sub>-C<sub>8</sub> linear or branched alkyl chains, methoxy, ethoxy, 2-hydroxyethoxy, 2-methoxyethoxy and 2-ethoxyethyl.

5 Representative but non-limiting examples of suitable monomers (c) are, for example, acrylic acid or methacrylic acid and salts, esters and amides thereof. The salts can be derived from any desired nontoxic metal, ammonium or substituted ammonium counterions.


10

The esters can be derived from C<sub>1</sub>-C<sub>40</sub> linear, C<sub>3</sub>-C<sub>40</sub> branched or C<sub>3</sub>-C<sub>40</sub> carbocyclic alcohols, from polyfunctional alcohols having from 2 to about 8 hydroxyl groups, such as ethylene glycol, hexylene glycol, glycerol and 1,2,6-hexanetriol, from

15 aminoalcohols or alcohol ethers such as methoxyethanol and ethoxyethanol, (alkyl)polyethylene glycols, (alkyl)polypropylene glycols or ethoxylated fatty alcohols, for example C<sub>12</sub>-C<sub>24</sub>-fatty alcohols reacted with 1 to 200 ethylene oxide units.

20 Also suitable are N,N-dialkylaminoalkyl acrylates and methacrylates and N-dialkylaminoalkylacryl- and -methacrylamides of the formula (III)

25



30

where

R<sup>17</sup> = H, alkyl having from 1 to 8 carbon atoms,

R<sup>18</sup> = H, methyl,

R<sup>19</sup> =        alkylene having from 1 to 24 carbon atoms, optionally substituted by alkyl,

35 substituted by alkyl,

$$R^{20}, R^{21} = C_1-C_{40} \text{ alkyl radical,}$$

$Z$  = nitrogen when  $g = 1$ , or oxygen when  $g = 0$ .

The amides can be unsubstituted, N-alkyl- or N-alkylamino-

40 monosubstituted or N,N-dialkyl-substituted or N,N-dialkylamino-disubstituted, where the alkyl or alkylamino groups are derived from C<sub>1</sub>-C<sub>40</sub> linear, C<sub>3</sub>-C<sub>40</sub> branched, or C<sub>3</sub>-C<sub>40</sub> carbocyclic units. In addition, the alkylamino groups can be quaternized.

45

## 11

Preferred comonomers of the formula III are

N,N-dimethylaminomethyl (meth)acrylate, N,N-diethylaminomethyl (meth)acrylate, N,N-dimethylaminoethyl (meth)acrylate, N,N-diethylaminoethyl (meth)acrylate,

- 5 N-[3-(dimethylamino)propyl]methacrylamide and  
N-[3-(dimethylamino)propyl]acrylamide.

Comonomers (c) which can likewise be used are substituted acrylic acids and salts, esters and amides thereof, where the


- 10 substituents on the carbon atoms are in the two or three position of the acrylic acid, and are independently of one another chosen from the group consisting of  $C_1$ - $C_4$ -alkyl, -CN, COOH particularly preferably methacrylic acid, ethacrylic acid and 3-cyanoacrylic acid. These salts, esters and amides of these substituted acrylic  
15 acids can be chosen as described above for the salts, esters and amides of acrylic acid.

Other suitable comonomers (c) are allyl esters of  $C_1$ - $C_{40}$  linear,  $C_3$ - $C_{40}$  branched or  $C_3$ - $C_{40}$  carbocyclic carboxylic acids, vinyl or  
20 allyl halides, preferably vinyl chloride and allyl chloride, vinyl ethers, preferably methyl, ethyl, butyl or dodecyl vinyl ether, vinylformamide, vinylmethylacetamide, vinylamine; vinyl lactams, preferably vinylpyrrolidone and vinylcaprolactam, vinyl- or allyl-substituted heterocyclic compounds, preferably

- 25 vinylpyridine, vinyloxazoline and allylpyridine.

Also suitable are N-vinylimidazoles of the formula IV, in which  $R^{22}$  to  $R^{24}$  independently of one another are hydrogen,  $C_1$ - $C_4$ -alkyl or phenyl:


30



35

Other suitable comonomers (c) are diallyl amines of the formula (V)

40



45

where  $R^{25} = C_1-$  to  $C_{24}$ -alkyl

- Other suitable comonomers (c) are vinylidene chloride; and hydrocarbons having at least one carbon-carbon double bond,
- 5 preferably styrene, alpha-methylstyrene, tert-butylstyrene, butadiene, isoprene, cyclohexadiene, ethylene, propylene, 1-butene, 2-butene, isobutylene, vinyltoluene, and mixtures of these monomers.
  - 10 Particularly suitable comonomers (c) are acrylic acid, methacrylic acid, ethyl acrylic acid, methyl acrylate, ethyl acrylate, propyl acrylate, n-butyl acrylate, iso-butyl acrylate, t-butyl acrylate, 2-ethylhexyl acrylate, decyl acrylate, methyl methacrylate, ethyl methacrylate, propyl methacrylate, n-butyl
  - 15 methacrylate, isobutyl methacrylate, t-butyl methacrylate, 2-ethylhexyl methacrylate, decyl methacrylate, methyl ethacrylate, ethyl ethacrylate, n-butyl ethacrylate, isobutyl ethacrylate, t-butyl ethacrylate, 2-ethylhexyl ethacrylate, decyl ethacrylate, stearyl (meth)acrylate, 2,3-dihydroxypropyl
  - 20 acrylate, 2,3-dihydroxypropyl methacrylate, 2-hydroxyethyl acrylate, hydroxypropyl acrylates, 2-hydroxyethyl methacrylate, 2-hydroxyethyl ethacrylate, 2-methoxyethyl acrylate, 2-methoxyethyl methacrylate, 2-methoxyethyl ethacrylate, 2-ethoxyethyl methacrylate, 2-ethoxyethyl ethacrylate,
  - 25 hydroxypropyl methacrylates, glyceryl monoacrylate, glyceryl monomethacrylate, polyalkylene glycol (meth)acrylates, unsaturated sulfonic acids such as, for example, acrylamidopropane sulfonic acid;
  - 30 acrylamide, methacrylamide, ethacrylamide, N-methylacrylamide, N,N-dimethylacrylamide, N-ethylacrylamide, N-isopropylacrylamide, N-butylacrylamide, N-t-butylacrylamide, N-octylacrylamide, N-t-octylacrylamide, N-octadecylacrylamide, N-phenylacrylamide, N-methylmethacrylamide, N-ethylmethacrylamide,
  - 35 N-dodecylmethacrylamide, 1-vinylimidazole, 1-vinyl-2-methylvinylimidazole, N,N-dimethylaminomethyl (meth)acrylate, N,N-diethylaminomethyl (meth)acrylate, N,N-dimethylaminoethyl (meth)acrylate, N,N-diethylaminoethyl (meth)acrylate, N,N-dimethylaminobutyl (meth)acrylate,
  - 40 N,N-diethylaminobutyl (meth)acrylate, N,N-dimethylaminohexyl (meth)acrylate, N,N-dimethylaminooctyl (meth)acrylate, N,N-dimethylaminododecyl (meth)acrylate, N-[3-(dimethylamino)propyl]methacrylamide, N-[3-(dimethylamino)propyl]acrylamide,
  - 45 N-[3-(dimethylamino)butyl]methacrylamide, N-[8-(dimethylamino)octyl]methacrylamide, N-[12-(dimethylamino)dodecyl]methacrylamide,

0050-0050-0050



N-[3-(diethylamino)propyl]methacrylamide,  
N-[3-(diethylamino)propyl]acrylamide;

maleic acid, fumaric acid, maleic anhydride and its half-esters,  
5 crotonic acid, itaconic acid, diallyldimethylammonium chloride,  
vinyl ethers (for example: methyl, ethyl, butyl or dodecyl vinyl  
ether), vinyl formamide, vinylmethylacetamide, vinylamine; methyl  
vinyl ketone, maleimide, vinylpyridine, vinylimidazole,  
vinylfuran, styrene, styrene sulfonate, allyl alcohol, and  
10 mixtures thereof.

Of these, particular preference is given to acrylic acid,  
methacrylic acid, maleic acid, fumaric acid, crotonic acid,  
maleic anhydride and its half-esters, methyl acrylate, methyl  
15 methacrylate, ethyl acrylate, ethyl methacrylate, n-butyl  
acrylate, n-butyl methacrylate, t-butyl acrylate, t-butyl  
methacrylate, isobutyl acrylate, isobutyl methacrylate,  
2-ethylhexyl acrylate, stearyl acrylate, stearyl methacrylate,  
N-t-butylacrylamide, N-octylacrylamide, 2-hydroxyethyl acrylate,  
20 hydroxypropyl acrylate, 2-hydroxyethyl methacrylate,  
hydroxypropyl methacrylate, alkylene glycol (meth)acrylates,  
styrene, unsaturated sulfonic acids such as, for example,  
acrylamidopropane sulfonic acid, vinylpyrrolidone,  
vinylcaprolactam, vinyl ethers (e.g.: methyl, ethyl, butyl or  
25 dodecyl vinyl ether), vinylformamide, vinylmethylacetamide,  
vinylamine, 1-vinylimidazole, 1-vinyl-2-methylimidazole,  
N,N-dimethylaminomethyl methacrylate and  
N-[3-(dimethylamino)propyl]methacrylamide;  
3-methyl-1-vinylimidazolium chloride, 3-methyl-1-vinylimidazolium  
30 methylsulfate, N,N-dimethylaminoethyl methacrylate,  
N-[3-(dimethylamino)propyl]methacrylamide quaternized with methyl  
chloride, methyl sulfate or diethyl sulfate.

Monomers having one basic nitrogen atom can be quaternized in the  
35 following manner:

Suitable for quaternizing the amines are, for example, alkyl  
halides having from 1 to 24 carbon atoms in the alkyl group, e.g.  
methyl chloride, methyl bromide, methyl iodide, ethyl chloride,  
40 ethyl bromide, propyl chloride, hexyl chloride, dodecyl chloride,  
lauryl chloride and benzyl halides, in particular benzyl chloride  
and benzyl bromide. Other suitable quaternizing agents are  
dialkyl sulfates, in particular dimethyl sulfate or diethyl  
sulfate. The quaternization of the basic amines can also be  
45 carried out with alkylene oxides such as ethylene oxide or

0050/49774



In some instances, it is also possible to use thio-containing silicone compounds.

Preference is given to using silicone-free regulators.

- 5 Crosslinking monomers which can be used are compounds having at least two ethylenically unsaturated double bonds, such as, for example, esters of ethylenically unsaturated carboxylic acids, such as acrylic acid or methacrylic acid and polyhydric alcohols, ethers of at least dihydric alcohols such as, for example, vinyl  
10 ethers or allyl ethers.

Examples of the parent alcohols are dihydric alcohols such as 1,2-ethanediol, 1,2-propanediol, 1,3-propanediol, 1,2-butanediol, 1,3-butanediol, 2,3-butanediol, 1,4-butanediol,

- 15 but-2-ene-1,4-diol, 1,2-pentanediol, 1,5-pentanediol, 1,2-hexanediol, 1,6-hexanediol, 1,10-decanediol, 1,2-dodecanediol, 1,12-dodecanediol, neopentyl glycol, 3-methylpentane-1,5-diol, 2,5-dimethyl-1,3-hexanediol, 2,2,4-trimethyl-1,3-pentanediol, 1,2-cyclohexanediol,  
20 1,4-cyclohexanediol, 1,4-bis(hydroxymethyl)cyclohexane, neopentyl glycol hydroxypivalate, 2,2-bis(4-hydroxyphenyl)propane, 2,2-bis[4-(2-hydroxypropyl)phenyl]propane, diethylene glycol, triethylene glycol, tetraethylene glycol, dipropylene glycol, tripropylene glycol, tetrapropylene glycol,  
25 3-thiopentane-1,5-diol, and polyethylene glycols, polypropylene glycols and polytetrahydrofurans having molecular weights of in each case 200 to 10 000. Apart from the homopolymers of ethylene oxide and propylene oxide, it is also possible to use block copolymers of ethylene oxide or propylene oxide or copolymers  
30 which contain ethylene oxide and propylene oxide groups in incorporated form. Examples of parent alcohols having more than two OH groups are trimethylolpropane, glycerol, pentaerythritol, 1,2,5-pentanetriol, 1,2,6-hexanetriol, triethoxycyanuric acid, sorbitan, sugars such as sucrose, glucose, mannose. It is of  
35 course also possible to use the polyhydric alcohols following reaction with ethylene oxide or propylene oxide, as the corresponding ethoxylates or propoxylates respectively. The polyhydric alcohols can also firstly be converted into the corresponding glycidyl ethers by reaction with epichlorohydrin.

- 40 Further suitable crosslinkers are the vinyl esters or the esters of monohydric, unsaturated alcohols with ethylenically unsaturated C<sub>3</sub>- to C<sub>6</sub>-carboxylic acids, for example acrylic acid, methacrylic acid, itaconic acid, maleic acid or fumaric acid.

- 45 Examples of such alcohols are allyl alcohol, 1-buten-3-ol, 5-hexen-1-ol, 1-octen-3-ol, 9-decen-1-ol, dicyclopentenyl alcohol, 10-undecen-1-ol, cinnamyl alcohol, citronellol, crotyl

alcohol or cis-9-octadecen-1-ol. However, it is also possible to esterify the monohydric, unsaturated alcohols with polybasic carboxylic acids, for example malonic acid, tartaric acid, trimellitic acid, phthalic acid, terephthalic acid, citric acid  
5 or succinic acid.

Further suitable crosslinkers are esters of unsaturated carboxylic acids with the above-described polyhydric alcohols, for example of oleic acid, crotonic acid, cinnamic acid or  
10 10-undecenoic acid.

Also suitable are straight-chain or branched, linear or cyclic aliphatic or aromatic hydrocarbons which have at least two double bonds which, in the case of aliphatic hydrocarbons, must not be  
15 conjugated, e.g., divinylbenzene, divinyltoluene, 1,7-octadiene, 1,9-decadiene, 4-vinyl-1-cyclohexene, trivinylcyclohexane or polybutadienes having molecular weights of from 200 to 20 000.

Also suitable are amides of unsaturated carboxylic acids, such  
20 as, for example, acrylic acid and methacrylic acid, itaconic acid, maleic acid and N-allylamines of at least difunctional amines, such as, for example, 1,2-diaminomethane, 1,2-diaminoethane, 1,3-diaminopropane, 1,4-diaminobutane, 1,6-diaminohexane, 1,12-dodecanediamine, piperazine,  
25 diethylenetriamine or isophorone diamine. Also suitable are the amides of allylamine and unsaturated carboxylic acids such as acrylic acid, methacrylic acid, itaconic acid, maleic acid, or at least dibasic carboxylic acids as have been described above.

30 Further suitable crosslinkers are triallylamine or corresponding ammonium salts, e.g. triallylmethylammonium chloride or triallylmethylammonium methyl sulfate.

It is also possible to use the N-vinyl compounds of urea  
35 derivatives, at least difunctional amides, cyanurates or urethanes, for example of urea, ethyleneurea, propyleneurea or tartramide, e.g. N,N'-divinylethyleneurea or N,N'-divinylpropyleneurea.

40 Further suitable crosslinkers are divinylldioxane, tetraallylsilane or tetravinylsilane.

Particularly preferred crosslinkers are, for example, methylenebisacrylamide, divinylbenzene, triallylamine and  
45 triallylammonium salts, divinylimidazole, N,N'-divinylethyleneurea, reaction products of polyhydric alcohols with acrylic acid or methacrylic acid, methacrylic esters and acrylic esters of

polyalkylene oxides or polyhydric alcohols which have been reacted with ethylene oxide and/or propylene oxide and/or epichlorohydrin, and allyl or vinyl ethers of polyhydric alcohols, for example 1,2-ethanediol, 1,4-butanediol, diethylene glycol, trimethylolpropane, glycerol, pentaerythritol, sorbitan and sugars such as sucrose, glucose, mannose.

- Very particularly preferred crosslinkers are pentaerythritol triallyl ethers, allyl ethers of sugars such as sucrose, glucose, mannose, divinylbenzene, methylenebisacrylamide, N,N'-divinylethylenurea, and (meth)acrylic esters of glycol, butanediol, trimethylolpropane or glycerol or (meth)acrylic esters of glycol, butanediol, trimethylolpropane or glycerol reacted with ethylene oxide and/or epichlorohydrin.
- The proportion of monomers which have a crosslinking action is 0 to 10% by weight, preferably 0.1 to 5% by weight, very particularly preferably 0.2 to 2% by weight.
- In the polymerization for the preparation of the polymers according to the invention, in some instances it is also possible for other polymers, such as, for example, polyamides, polyurethanes, polyesters, homo- and copolymers of ethylenically unsaturated monomers, to be present. Examples of such polymers, some of which are also used in cosmetics, are the polymers known under the tradenames Amerhold™, Ultrahold™, Ultrahold Strong™, Luviflex™ VBM, Luvimer™, Acronal™, Acudyne™, Stepanhold™, Lovocryl™, Versatyl™, Amphomer™ or Eastma AQ™.
- The comonomers (c) according to the invention can, provided they contain ionizable groups, be partially or completely neutralized using acids or bases before or after the polymerization in order, for example, to adjust the solubility or dispersibility in water to a desired degree.
- Neutralizing agents for monomers carrying acid groups which can be used are, for example, mineral bases such as sodium carbonate, alkali metal hydroxides and ammonia, organic bases such as aminoalcohols, specifically 2-amino-2-methyl-1-propanol, monoethanolamine, diethanolamine, triethanolamine, triisopropanolamine, tri[(2-hydroxy)1-propyl]amine, 2-amino-2-methyl-1,3-propanediol, 2-amino-2-hydroxymethyl-1,3-propanediol and diamines, such as, for example, lysine.
- To prepare the polymers, the monomers of component a) can be polymerized in the presence of the polyethers either using initiators which form free radicals, or by the action of

high-energy radiation, which is also intended to mean the action of high-energy electrons.

Initiators which can be used for the free-radical polymerization are the peroxo and/or azo compounds customary for this purpose, for example alkali metal or ammonium peroxydisulfates, diacetyl peroxide, dibenzoyl peroxide, succinyl peroxide, di-tert-butyl peroxide, tert-butyl perbenzoate, tert-butyl perpivalate, tert-butyl peroxy-2-ethylhexanoate, tert-butyl permaleate, cumene hydroperoxide, diisopropyl peroxydicarbamate, bis-(o-toluoyl) peroxide, didecanoyl peroxide, dioctanoyl peroxide, dilauroyl peroxide, tert-butyl perisobutyrate, tert-butyl peracetate, di-tert-amyl peroxide, tert-butyl hydroperoxide, azobisisobutyronitrile, azobis-(2-amidinopropane) dihydrochloride or 2,2'-azobis(2-methyl-butyronitrile). Also suitable are initiator mixtures or redox initiator systems, such as, for example, ascorbic acid/iron(II) sulfate/sodium peroxodisulfate, tert-butyl hydroperoxide/sodium disulfite, tert-butyl hydroperoxide/sodium hydroxymethanesulfinate.

Preference is given to using organic peroxides.

The amounts of initiator or initiator mixtures used, based on monomer used, are between 0.01 and 10% by weight, preferably between 0.1 and 5% by weight.

The polymerization is carried out in a temperature range from 40 to 200°C, preferably in the range from 50 to 140°C, particularly preferably in the range from 60 to 110°C. It is usually carried out under atmospheric pressure, but can also be carried out under reduced or increased pressure, preferably between 1 and 5 bar.

The polymerization can, for example, be carried out as solution polymerization, bulk polymerization, emulsion polymerization, inverse emulsion polymerization, suspension polymerization, inverse suspension polymerization or precipitation polymerization, without the possible methods being limited thereto.

In the case of bulk polymerization, the procedure may involve dissolving the polyether-containing compound b) in at least one monomer of group a) and possibly other comonomers of group c) and, after the addition of a polymerization initiator, fully polymerizing the mixture. The polymerization can also be carried out semicontinuously by firstly introducing some, e.g. 10%, of the mixture to be polymerized comprising the polyether-containing compound b), at least one monomer from group a), possibly other comonomers of group c) and initiator, heating the mixture to the

polymerization temperature and after the polymerization has started, adding the remainder of the mixture to be polymerized in accordance with the progress of the polymerization. The polymers can also be obtained by initially introducing the

5 polyether-containing compounds of group b) into a reactor, heating them to the polymerization temperature and adding at least one monomer of group a), possibly other comonomers of group c) and polymerization initiator either in one portion, step by step or, preferably, continuously, and polymerizing.

10

If desired, the above described polymerization can also be carried out in a solvent. Suitable solvents are, for example, alcohols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol, sec-butanol, tert-butanol, n-hexanol and cyclohexanol,

15 and glycols, such as ethylene glycol, propylene glycol and butylene glycol, and the methyl or ethyl ethers of dihydric alcohols, diethylene glycol, triethylene glycol, glycerol and dioxane. The polymerization can also be carried out in water as solvent. In this case, the initial batch is a solution which,  
20 depending on the amount of monomers of component a) added, is soluble in water to a greater or lesser degree. In order to convert water-insoluble products, which can form during the polymerization, into solution, it is possible, for example, to add organic solvents, such as monohydric alcohols having from 1  
25 to 3 carbon atoms, acetone or dimethylformamide. However, in the case of polymerization in water, it is also possible to convert the water-insoluble polymers into a finely divided dispersion by addition of customary emulsifiers or protective colloids, e.g. polyvinyl alcohol.

30

The emulsifiers used are, for example, ionic or nonionic surfactants whose HLB value is in the range from 3 to 13. The definition of the HLB value can be found in the publication by W.C. Griffin, J. Soc. Cosmetic Chem., Volume 5, 249 (1954).

35

The amount of surfactants, based on the polymer, is from 0.1 to 10% by weight. Using water as solvent gives solutions or dispersions of the polymers. If solutions of the polymer are prepared in an organic solvent or in mixtures of an organic  
40 solvent and water, then, per 100 parts by weight of the polymer, from 5 to 2000, preferably from 10 to 500, parts by weight of the organic solvent or of the solvent mixture are used.

Preference is given to polymers obtainable by free-radical  
45 polymerization of

## 20

- a) 10 - 98 % by weight of at least one vinyl ester of  $C_1-C_{24}$  carboxylic acids in the presence of
- b) 2 - 90 % by weight of at least one polyether-containing compound and
- 5 c) 0 - 50 % by weight of one or more other copolymerizable monomers.

Particular preference is given to polymers obtainable by free-radical polymerization of

10

- a) 50 - 97 % by weight of at least one vinyl ester of  $C_1-C_{24}$  carboxylic acids in the presence of
- b) 3 - 50 % by weight of at least one polyether-containing compound and
- 15 c) 0 - 30 % by weight of one or more other copolymerizable monomers.

Very particular preference is given to polymers obtainable by free-radical polymerization of

20

- a) 60 - 97 % by weight of at least one vinyl ester of  $C_1-C_{24}$  carboxylic acids in the presence of
- b) 3 - 40 % by weight of at least one polyether-containing compound and
- 25 c) 0 - 20 % by weight of one or more other copolymerizable monomers.


To prepare the polymers used according to the invention, the ester groups of the original monomers a) and optionally of other  
30 monomers are cleaved after the polymerization by hydrolysis, alcoholysis, or aminolysis. This process step is generally referred to below as hydrolysis. The hydrolysis takes place in a manner known per se by the addition of a base, preferably the addition of a sodium or potassium hydroxide solution in water  
35 and/or alcohol. Particular preference is given to using methanolic sodium or potassium hydroxide solutions. The hydrolysis is carried out at temperatures in the range from 10 to 80°C, preferably in the range from 20 to 60°C. The degree of hydrolysis depends on the amount of base used, on the hydrolysis  
40 temperature, on the hydrolysis time and the water content of the solution.

The degree of hydrolysis of the polyvinyl ester groups is in the range from 1 to 100%, preferably in the range from 40 to 100%,  
45 particularly preferably in the range from 65 to 100%, very particularly preferably in the range from 80 to 100%.



The polymers prepared in this way can then be cationized by reaction of hydroxyl and/or amino functions present in the polymer with epoxides of the formula VI ( $R^{26} = C_1-$  to  $C_{40}$ -alkyl).

5



- 10 For this, the hydroxyl groups of the polyvinyl alcohol units and vinylamine units, formed by hydrolysis of vinylformamide, can preferably be reacted with the epoxides.

The epoxides of the formula VI can also be produced in situ by reaction of the corresponding chlorohydrins with bases, for

- 15 example sodium hydroxide.

Preference is given to using 2,3-epoxypropyltrimethylammonium chloride or 3-chloro-2-hydroxypropyltrimethylammonium chloride. The K values of the polymers should be in the range from 10 to 300, preferably 25 to 250, particularly preferably 25 to 200,

- 20 very particularly preferably in the range from 30 to 150. The K value desired in each case can be adjusted in a manner known per se through the composition of the feed substances. The K values are determined in accordance with Fikentscher, Cellulosechemie, Vol. 13, p. 58 to 64, and 71 to 74 (1932) in N-methylpyrrolidone  
25 at 25°C and polymer concentrations which, depending on the K value range, are between 0.1 % by weight and 5 % by weight.

After the hydrolysis, the polymer solutions can be steam distilled to remove any solvents. After the steam distillation,

- 30 aqueous solutions or dispersions are obtained depending on the degree of hydrolysis, type of polyethers b), of vinyl esters a) and any monomers c) used.

The polymers obtained can also be subsequently crosslinked by

- 35 reacting the hydroxyl groups or amino groups in the polymer with at least bifunctional reagents. In the case of low degrees of crosslinking, water-soluble products are obtained, and in the case of high degrees of crosslinking, water-swellaable or insoluble products are obtained.

40

For example, the polymers according to the invention can be reacted with aldehydes, dialdehydes, ketones and diketones, e.g. formaldehydes, acetaldehyde, glyoxal, glutaraldehyde, succindialdehyde or terephthalaldehyde. Also suitable are

- 45 aliphatic or aromatic carboxylic acids, for example maleic acid, oxalic acid, malonic acid, succinic acid or citric acid, or carboxylic acid derivatives, such as carboxylic esters,

anhydrides or halides. Also suitable are polyfunctional epoxides, e.g. epichlorohydrin, glycidyl methacrylate, ethylene glycol diglycidyl ether, 1,4-butanediol diglycidyl ether or 1,4-bis(glycidyloxy)benzene. Also suitable are diisocyanates, for example hexamethylene diisocyanate, isophorone diisocyanate, methylenediphenyl diisocyanate, toluylene diisocyanate or divinylsulfone.

Also suitable are inorganic compounds, such as boric acid or boric acid salts, which are referred to collectively as borates below, for example sodium metaborate, borax (disodium tetraborate), and salts of polyvalent cations, e.g. copper(II) salts, such as copper(II) acetate or zinc, aluminum, titanium salts.

Boric acid and/or boric acid salts, such as sodium metaborate or disodium tetraborate, are preferably suitable for the subsequent crosslinking. In this connection, boric acid and/or boric acid salts can, preferably as salt solutions, be added to the solutions of the polymers according to the invention. Preference is given to adding boric acid and/or boric acid salts to the aqueous polymer solutions.

The boric acid and/or boric acid salts can be added to the polymer solutions directly after preparation. It is, however, also possible to add the boric acid and/or boric acid salts subsequently to the cosmetic formulations containing the polymers according to the invention, or to add them during the preparation process of the cosmetic formulations.

The proportion of boric acid and/or boric acid salts, based on the polymers according to the invention, is 0 to 15% by weight, preferably 0 to 10% by weight, particularly preferably 0 to 5% by weight.

The polymer solutions and dispersions can be converted into powder form by a variety of drying methods, such as, for example, spray drying, fluidized spray drying, drum drying or freeze drying. The drying method used in preference is spray drying. The dry polymer powder obtained in this way can be used to prepare an aqueous solution or dispersion again, by dissolution or redispersion in water. Conversion into powder form has the advantage of better storability, easier transportation, and a lower propensity for microbial attack.

5 The water-soluble or water-dispersible polyalkylene oxide- or polyglycerol-containing polymers according to the invention are highly suitable for use in hair cosmetic formulations.

The polymers according to the invention, prepared by free-radical polymerization of vinyl esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters, are suitable as styling agents and/or conditioning agents in hair cosmetic preparations such as hair treatments, hair lotions, hair rinses, hair emulsions, fluids for treating hair ends, neutralizing agents for permanent waves, "hot-oil-treatment" preparations, conditioners, setting lotions or hairsprays. Depending on the area of application, the hair cosmetic preparations can be applied as a spray, foam, gel, gel spray or mousse.

The hair cosmetic formulations according to the invention comprise, in a preferred embodiment,

- 25 a) 0.05 - 20 % by weight of the polymer according to the invention, prepared by free-radical polymerization of vinyl esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at least partial hydrolysis of the ester functions of the
- 30 original vinyl esters
- b) 20 - 99.95 % by weight of water and/or alcohol
- c) 0 - 79.5 % by weight of other constituents

Alcohol is taken to mean all alcohols customary in cosmetics,  
35 e.g. ethanol, isopropanol, n-propanol.

Other constituents are taken to mean the additives customary in cosmetics, for example propellants, antifoams, surface-active compounds, i.e. surfactants, emulsifiers, foam formers and  
40 solubilizers. The surface-active compounds used can be anionic, cationic, amphoteric or neutral. In addition, other customary constituents can be, for example, preservatives, perfume oils, opacifiers, active ingredients, UV filters, care substances such as panthenol, collagen, vitamins, protein hydrolysates, alpha-  
45 and beta-hydroxycarboxylic acids, stabilizers, pH regulators,

dyes, viscosity regulators, gel formers, salts, humectants, refatting agents and other customary additives.

These also include all styling and conditioning polymers known in cosmetics which can be used in combination with the polymers according to the invention, in cases where very specific properties are to be set.

- Suitable traditional hair cosmetic polymers are, for example,
- 10 anionic polymers. Such anionic polymers are homo- and copolymers of acrylic acid and methacrylic acid or salts thereof, copolymers of acrylic acid and acrylamide and salts thereof; sodium salts of polyhydroxycarboxylic acids, water-soluble or water-dispersible polyesters, polyurethanes (Luviset® P.U.R.) and polyureas.
  - 15 Particularly suitable polymers are copolymers of t-butyl acrylate, ethyl acrylate, methacrylic acid (e.g. Luvimer® 100P), copolymers of N-tert-butylacrylamide, ethyl acrylate, acrylic acid (Ultrahold® 8, strong), copolymers of vinyl acetate, crotonic acid and optionally other vinyl esters (e.g. Luviset® grades),
  - 20 maleic anhydride copolymers, optionally reacted with alcohols, anionic polysiloxanes, e.g. carboxy-functional ones, copolymers of vinylpyrrolidone, t-butyl acrylate, methacrylic acid (e.g. Luviskol® VBM).
  - 25 Very particularly preferred anionic polymers are acrylates with an acid number greater than or equal to 120 and copolymers of t-butyl acrylate, ethyl acrylate and methacrylic acid.

- Other suitable hair cosmetic polymers are cationic polymers with
- 30 the name polyquaternium according to INCI, e.g. copolymers of vinylpyrrolidone/N-vinylimidazolium salts (Luviquat® FC, Luviquat® HM, Luviquat® MS, Luviquat® Care), copolymers of N-vinylpyrrolidone/dimethylaminoethyl methacrylate, quaternized with diethylsulfate (Luviquat® PQ 11), copolymers of
  - 35 N-vinylcaprolactam-N-vinylpyrrolidone/N-vinylimidazolium salts (Luviquat® Hold); cationic cellulose derivatives (polyquaternium-4 and -10), acrylamide copolymers (polyquaternium-7).

- Other suitable hair cosmetic polymers are also neutral polymers
- 40 such as polyvinylpyrrolidones, copolymers of N-vinylpyrrolidone and vinyl acetate and/or vinyl propionate, polysiloxanes, polyvinylcaprolactam and copolymers with N-vinylpyrrolidone, polyethyleneimines and salts thereof, polyvinylamines and salts thereof, cellulose derivatives, polyaspartic acid salts and
  - 45 derivatives.

To establish certain properties, the preparations can also additionally comprise conditioning substances based on silicone compounds. Suitable silicone compounds are, for example, polyalkylsiloxanes, polyarylsiloxanes, polyaryalkylsiloxanes, 5 polyethersiloxanes, silicone resins or dimethicone copolyols (CTFA) and amino-functional silicone compounds such as Amodimethicones (CTFA).

The polymers according to the invention are particularly suitable 10 as setting agents in hair styling preparations, in particular hairsprays (aerosols and pump sprays without propellant) and hair foams (aerosol foams and pump foams without propellant). In a preferred embodiment, these preparations comprise

- 15 a) 0.1 - 10 % by weight of the polymer according to the invention, prepared by free-radical polymerization of vinyl esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at least partial hydrolysis of the ester functions of the
- 20 original vinyl esters
- b) 20 - 99.9 % by weight of water and/or alcohol
- c) 0 - 70 % by weight of a propellant
- d) 0 - 20 % by weight of other constituents.
- 25 The propellants are those customarily used for hairsprays or aerosol foams. Preference is given to mixtures of propane/butane, pentane, dimethyl ether, 1,1-difluoroethane (HFC-152 a), carbon dioxide, nitrogen or compressed air.
- 30 A formulation preferred according to the invention for aerosol hair foams comprises
- a) 0.1 - 10 % by weight of the polymer according to the invention, prepared by free-radical polymerization of vinyl
- 35 esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters
- b) 55 - 94.8 % by weight of water and/or alcohol
- 40 c) 5 - 20 % by weight of a propellant
- d) 0.1 - 5 % by weight of an emulsifier
- e) 0 - 10 % by weight of other constituents.

The emulsifiers can be any emulsifiers customarily used in hair 45 foams. Suitable emulsifiers can be nonionic, cationic or anionic.

0050/49774

Examples of nonionic emulsifiers (INCI nomenclature) are laureths, e.g. laureth-4; ceteths, e.g. cetheth-1, polyethylene glycol cetyl ether; cetareths, e.g. cetheareth-25, polyglycol fatty acid glycerides, hydroxylated lecithin, lactyl esters of

5 fatty acids, alkylpolyglycosides.

Examples of cationic emulsifiers are cetyldimethyl-2-hydroxy-ethylammonium dihydrogenphosphate, cetyltrimonium chloride, cetyltrimonium bromide, cocotrimonium methylsulfate, quaternium-1 to x (INCI).

- 10 Anionic emulsifiers can, for example, be chosen from the group consisting of alkylsulfates, alkyl ether sulfates, alkylsulfonates, alkylarylsulfonates, alkylsuccinates, alkylsulfosuccinates, N-alkoylsarcosinates, acyltaurates, acylisethionates, alkylphosphates, alkyl ether phosphates, alkyl
- 15 ether carboxylates, alpha-olefin sulfonates, in particular the alkali metal and alkaline earth metal salts, e.g. sodium, potassium, magnesium, calcium, and ammonium and triethanolamine salts. The alkyl ether sulfates, alkyl ether phosphates and alkyl
- 20 ether carboxylates can have between 1 and 10 ethylene oxide or propylene oxide units, preferably from 1 to 3 ethylene oxide units, in the molecule.

A preparation which is suitable according to the invention for styling gels can, for example, have the following composition:

- 25 a) 0.1 - 10 % by weight of the polymer according to the invention, prepared by free-radical polymerization of vinyl esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at
- 30 least partial hydrolysis of the ester functions of the original vinyl esters
- b) 60 - 99.85 % by weight of water and/or alcohol
- c) 0.05 - 10 % by weight of a gel former
- d) 0 - 20 % by weight of other constituents.

- 35 The gel formers which can be used are any gel formers customary in cosmetics. These include totally crosslinked polyacrylic acid, for example carbomer (INCI), cellulose derivatives, e.g. hydroxypropylcellulose, hydroxyethylcellulose, cationically
- 40 modified celluloses, polysaccharides, e.g. xanthan gum, caprylic/capric triglycerides, sodium acrylates copolymer, polyquaternium-32 (and) paraffinum liquidum (INCI), sodium acrylates copolymer (and) paraffinum liquidum (and) PPG-1 trideceth-6, acrylamidopropyltrimonium chloride/acrylamide
- 45 copolymer, steareth-10 allyl ether acrylates copolymer, polyquaternium-37 (and) paraffinum liquidum (and) PPG-1 trideceth-6, polyquaternium 37 (and) propylene glycole dicaprate

dicaprylate (and) PPG-1 trideceth-6, polyquaternium-7, polyquaternium-44.

The polymers according to the invention can also be used in  
5 shampoo formulations as setting and/or conditioning agents. Suitable conditioning agents are, in particular, polymers with a cationic charge.

Preferred shampoo formulations comprise

- 10 a) 0.05 - 10 % by weight of the polymer according to the invention, prepared by free-radical polymerization of vinyl esters and optionally other polymerizable monomers in the presence of polyether-containing compounds and subsequent at least partial hydrolysis of the ester functions of the  
15 original vinyl esters
- b) 25 - 94.95 % by weight of water
- c) 5 - 50 % by weight of surfactants
- c) 0 - 5 % by weight of another conditioning agent
- d) 0 - 10 % by weight of other cosmetic constituents.

20 In the shampoo formulations, it is possible to use all anionic, neutral, amphoteric or cationic surfactants used customarily in shampoos.

- 25 Suitable anionic surfactants are, for example, alkylsulfates, alkyl ether sulfates, alkylsulfonates, alkylarylsulfonates, alkylsuccinates, alkylsulfo succinates, N-alkoylsarcosinates, acyltaurates, acylisethionates, alkylphosphates, alkyl ether phosphates, alkyl ether carboxylates, alpha-olefinsulfonates, in  
30 particular the alkali metal and alkaline earth metal salts, e.g. sodium, potassium, magnesium, calcium, and ammonium and triethanolamine salts. The alkyl ether sulfates, alkyl ether phosphates and alkyl ether carboxylates can have between 1 and 10 ethylene oxide or propylene oxide units, preferably from 1 to 3  
35 ethylene oxide units, in the molecule.

Suitable examples are sodium laurylsulfate, ammonium laurylsulfate, sodium lauryl ether sulfate, ammonium lauryl ether sulfate, sodium lauroyl sarcosinate, sodium oleyl succinate,  
40 ammonium lauryl sulfo succinate, sodium dodecylbenzenesulfonate, triethanolamine dodecylbenzenesulfonate.

Suitable amphoteric surfactants are, for example, alkylbetaines, alkylamidopropylbetaines, alkylsulfo betaines, alkylglycinates,  
45 alkylcarboxyglycinates, alkylamphoacetates or -propionates, alkylamphodiacetates or -dipropionates.

It is, for example, possible to use cocodimethylsulfopropylbetaine, laurylbetaine, cocamidopropylbetaine or sodium cocamphopropionate.

- 5 Examples of suitable nonionic surfactants are the reaction products of aliphatic alcohols or alkyl phenols having from 6 to 20 carbon atoms in the alkyl chain, which can be linear or branched, with ethylene oxide and/or propylene oxide. The amount of alkylene oxide is from about 6 to 60 mole per mole of alcohol.
- 10 Also suitable are alkylamine oxides, mono- or dialkyl alkanolamides, fatty acid esters of polyethylene glycols, alkyl polyglycosides or sorbitan ether esters.

The shampoo formulations can also comprise customary cationic

15 surfactants, such as, for example, quaternary ammonium compounds, for example cetyltrimethylammonium chloride.

- To achieve certain effects, customary conditioning agents can be used in combination with the polymers according to the invention
- 20 in the shampoo formulations. Such conditioning agents include, for example, cationic polymers with the name polyquaternium according to INCI, in particular copolymers of vinylpyrrolidone/N-vinylimidazolium salts (Luviquat® FC, Luviquat® HM, Luviquat® MS, Luviquat® Care), copolymers of N-vinylpyrrolidone/dimethyl-
- 25 aminoethyl methacrylate, quaternized with diethyl sulfate (Luviquat® PQ 11), copolymers of N-vinylcaprolactam/N-vinylpyrrolidone/N-vinylimidazolium salts (Luviquat® Hold); cationic cellulose derivatives (polyquaternium-4 and -10), acrylamide copolymers (polyquaternium-7). In addition, it is also possible
- 30 to use protein hydrolysates, and conditioning substances based on silicone compounds, for example polyalkylsiloxanes, polyarylsiloxanes, polyarylalkylsiloxanes, polyether siloxanes or silicone resins. Other suitable silicone compounds are Dimethicone Copolyols (CTFA) and amino-functional silicone
- 35 compounds such as Amodimethicones (CTFA).


The invention further provides polymers obtainable by free-radical polymerization of

- 40 a) at least one vinyl ester of a C<sub>1</sub>-C<sub>24</sub>-carboxylic acid, in the presence of
- b) polyether-containing silicone derivatives and
- c) optionally one or more other copolymerizable monomers
- 45 and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters.







Preference is given to polymers obtainable by free-radical polymerization of

- a) at least one vinyl ester of a C<sub>1</sub>-C<sub>24</sub>-carboxylic acid, in the presence of
- b) polyether-containing silicone derivatives which contain the following structural elements:




where:



- R<sup>13</sup> is a C<sub>1</sub>-40 organic radical which can contain amino, carboxyl or sulfonate groups, or where e = 0, is also the anion of an inorganic acid,

- and where the radicals R<sup>8</sup> can be identical or different, and come either from the group of aliphatic hydrocarbons having from 1 to 20 carbon atoms, are cyclic aliphatic hydrocarbons having from 3 to 20 carbon atoms, are of an aromatic nature or are identical to R<sup>12</sup>, where:



5

with the proviso that at least one of the radicals  $R^8$ ,  $R^9$  or  $R^{10}$  is a polyalkylene oxide-containing radical as defined above,

10 and  $f$  is an integer from 1 to 6,

$a$  and  $b$  are integers such that the molecular weight of the polysiloxane block is between 300 and 30000,

$c$  and  $d$  can be integers between 0 and 50, with the proviso that the sum  $c + d$  is greater than 0, and  $e$  is 0 or 1,

15

and

c) optionally one or more other copolymerizable monomers


20 and subsequent at least partial hydrolysis of the vinyl ester functions of the original vinyl esters.

Very particular preference is given to polymers obtainable by free-radical polymerization of

25 a) at least one vinyl ester of a  $C_1$ - $C_{24}$ -carboxylic acid, in the presence of

b) polyether-containing silicone derivatives of the structure:

30



35

and

c) optionally one or more other copolymerizable monomers

40

and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters a).

The invention further provides polymers obtainable by

45 free-radical polymerization of

a) a vinyl ester of a  $C_1$ - $C_{24}$  carboxylic acid in the presence of

## 31

- b) polyether-containing compounds obtainable by reaction of polyethyleneimines with alkylene oxides and
- c) optionally one or more other copolymerizable monomers

5 and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters.

The invention further provides polymers obtainable by free-radical polymerization of

10

- a) a vinyl ester of a  $C_1$ - $C_{24}$  carboxylic acid in the presence of
- b) homo- and copolymers of ethylenically unsaturated polyether-containing compounds and
- c) optionally one or more other copolymerizable monomers

15

and subsequent at least partial hydrolysis of the ester functions of the original vinyl esters.

The invention further provides crosslinked polymeres obtainable

20 by free-radical polymerization of

- a) at least one vinyl ester of  $C_1$ - $C_{24}$  carboxylic acids in the presence of
- b) polyether-containing compounds and
- 25 c) optionally one or more further copolymerizable monomers

and subsequent at least partial hydrolysis of the ester functions of the original monomers a), where the crosslinker used is either already present during the polymerization, or is added after the  
30 polymerization and hydrolysis.

Preparation examples:

Preparation procedure for Examples 1 to 26

35

The polyether-containing compound is heated to 80°C in a polymerization vessel with stirring and under a gentle stream of nitrogen. With stirring, vinyl acetate and optionally the other monomers are metered in over the course of 3 h. At the same time,  
40 a solution of 1.4 g of tert-butyl perpivalate in 30 g of methanol is likewise added over 3 h. The mixture is then stirred for a further 2 h at 80°C. After cooling, the polymer is dissolved in 450 ml of methanol. For hydrolysis, 50 ml of a 10% strength methanolic sodium hydroxide solution are added at 30°C. After  
45 about 40 min., the reaction is terminated by addition of 750 ml

of 1% strength acetic acid. The methanol is removed by distillation.

The K values were determined at a concentration of 1% in 5 N-methylpyrrolidone.

Table

	Exam- ple	Graft base	Vinyl ester	Comonomer	K value	Degree of hydrolysis [%]
10	1	PEG 1500 <sup>1</sup> , 72 g	vinyl acetate, 410 g	-	47	> 95
	2	PEG 4000, 72 g	vinyl acetate, 410 g	-	51	> 95
15	3	PEG 6000, 72 g	vinyl acetate, 410 g	-	54	> 95
	4	PEG 6000, 137 g	vinyl acetate, 410 g		49	> 95
	5	PEG 6000, 22 g	vinyl acetate, 410 g	-	73	> 95
20	6	PEG 6000, 410 g	vinyl acetate, 410 g		42	> 95
	7	PEG 9000, 137 g	vinyl acetate, 410 g	-	58	> 95
	8	polyglycerol 2200, 72 g	vinyl acetate, 410 g	-	66	> 95
25	9	PEG-PPG block copolymer 8000 <sup>2</sup> , 72 g	vinyl acetate, 410 g	-	45	> 95
	10	methylpolyethylene glycol 2000 <sup>3</sup> , 72 g	vinyl acetate, 410 g	-	47	> 95
30	11	alkylpolyethylene glycol 3500 <sup>4</sup> , 72 g	vinyl acetate, 410 g	-	48	> 95
	12	PPG 4000 <sup>5</sup> , 72 g	vinyl acetate, 410 g		50	> 95
35	13	PEG 20000, 72 g	vinyl acetate, 410 g	-	69	> 95
	14	PEG 20000, 103 g	vinyl acetate, 410 g	-	64	> 95
	15	PEG 20000, 137 g	vinyl acetate, 410 g	-	59	> 95
40	16	PEG 20000, 615 g	vinyl acetate, 410 g	-	55	86
	17	PEG 35000, 72 g	vinyl acetate, 410 g	-	77	> 95
45	18	PEG 35000, 137 g	vinyl acetate, 410 g	-	80	> 95
	19	PEG 35000, 205 g	vinyl acetate, 410 g	-	65	97

Example	Graft base	Vinyl ester	Comonomer	K value	Degree of hydrolysis [%]
20	dimethicone copolyol <sup>6</sup> , 202 g	vinyl acetate, 410 g	-	58	> 95
21	poly(sodium methacrylate-co-methylpolyethylene glycol methacrylate) <sup>7</sup> 103 g	vinyl acetate, 410 g		43	> 95
22	ethoxylated polyethyleneimine <sup>8</sup>	vinyl acetate, 410 g		52	> 95
23	PEG 6000, 72 g	vinyl acetate, 386 g	methyl methacrylate, 24 g	47	> 95
24	PEG 20000, 72 g	vinyl acetate, 328 g	N-vinylpyrrolidone, 82 g	61	> 95
25	PEG 20000, 72 g	vinyl acetate, 362 g	3-methyl-1-vinylimidazolium methylsulfate, 48 g	53	> 95
26	PEG 6000, 72 g	vinyl acetate, 367 g	N-vinylformamide, 41 g	57	> 95
27	PEG 6000, 72 g	vinyl acetate, 326 g	N-vinylformamide, 82 g	67	> 95
28	PEG 35000, 270 g	vinyl acetate, 410 g		59	96
29	PEG 35000, 270 g	vinyl acetate, 410 g	pentaerythritol triallyl ether, 1.6 g	71	95
30	PEG 35000, 270 g	vinyl acetate, 410 g	pentaerythritol triallyl ether, 0.8 g	65	94
31	PEG 35000, 270 g	vinyl acetate, 410 g	N,N'-divinylethyleneurea, 0.7 g	73	95
32	PEG 12000, 270 g	vinyl acetate, 410 g	pentaerythritol triallyl ether, 1.6 g	50	94

- 1 PEG x: polyethylene glycol with an average molecular weight x (weight average)  
 2 Lutrol F 68 from BASF Aktiengesellschaft (PPG: polypropylene glycol)  
 3 Pluriol A 2000 E from BASF Aktiengesellschaft  
 4 Lutensol AT 80 from BASF Aktiengesellschaft (C<sub>16</sub>-C<sub>18</sub> fatty alcohol + 80 EO)  
 5 Polypropylene glycol with an average molecular weight of 4000  
 6 Belsil DMC 6031TM from Wacker Chemie GmbH  
 7 Molar ratio of sodium methacrylate/methylpolyethylene glycol methacrylate 4:1, methylpolyethylene glycol with a molar mass of about 1000  
 8 Prepared from 12.5% of polyethyleneimine (average molecular weight 1400) and 87.5% of ethylene oxide

## 34

Example 33: Reaction with 3-chloro-2-hydroxypropyltrimethylammonium chloride

22 g of a 60% strength aqueous solution of

- 5 3-chloro-2-hydroxypropyltrimethylammonium chloride and 3.5 g of sodium hydroxide are added to 400 g of a 32.9% strength solution from Example 3. The mixture is stirred for 3 hours at 60°C and then for a further two hours at 90°C, giving a clear solution.

- 10 Example 34: Reaction with 3-chloro-2-hydroxypropyltrimethylammonium chloride

46 g of a 60% strength aqueous solution of

- 15 3-chloro-2-hydroxypropyltrimethylammonium chloride and 6 g of sodium hydroxide are added to 400 g of a 15.3% strength solution from Example 26. The mixture is stirred for 3 hours at 60°C and then for a further two hours at 90°C, giving a clear solution.

Example 35:

20

At room temperature and with stirring, a 5% strength aqueous solution of disodium tetraborate decahydrate (borax) is added to a 19.3% strength aqueous solution of the polymer from example 28 over the course of half an hour. An increase in the viscosity is

25 observed.

Amount of 5% strength borax solution added [g]	Brookfield viscosity (LVF, spindle 2, 30 rpm, 23°C) [mPas]
0	110
14.9	128
18.0	216
21.0	534
24.0	2 228
26.9	7 520 <sup>1</sup>
29.8	29 190 <sup>2</sup>

35

1 Spindle 4, 30 rpm

2 Spindle 4, 6 rpm

Formulation Examples:

40

Example 36:

Aerosol hair foam formulation:

2.00 % of copolymer from Example 3

- 45 2.00 % of Luviquat Mono LS (cocotrimonium methylsulfate)

67.7 % of water

10.0 % of propane/butane 3.5 bar (20°C)

q.s. perfume oil

Example 37 (Comparative Example):

- 5 2.00 % polymer content Luviquat Hold (polyquaternium-46)  
 2.00 % of Luviquat Mono LS (cocotrimonium methylsulfate)  
 67.7 % of water  
 10.0 % of propane/butane 3.5 bar (20°C)  
 q.s. perfume oil

10

Using Example 36 and Example 37 (Comparative Example), half-head tests were carried out on dummy heads. The assessment was subjectively carried out by trained hairdressers and laboratory assistants.

15

Scale of grades from 1 (very good) to 3 (poor)

	Example 36	Example 37 (Comparative Example)
	1	1
20	1	1
	1	1
	1-	2
	1-	2+
	1	2+
25	2+	2
	1	1-
	1-	2+
	1	2-

- 30 The formulation from Example 36 had better hold, better wet combability, lower tack and better hair elasticity than the formulation from Example 37 (Comparative Example).

Example 38:

35 Aerosol hair foam:

INCI

4.00 % of copolymer from Example 19

0.20 % of Cremophor A 25

Ceteareth-25

1.00 % of Luviquat Mono CP

Hydroxyethyl cetyldimonium  
phosphate

40

5.00 % of ethanol

1.00 % of Panthenol

10.0 % of propane/butane 3.5 bar (20°C)

q.s. perfume oil

45 ad 100 % with water

Example 39:

Pump foam:

- 2.00 % of copolymer from Example 7  
5 2.00 % of Luviflex Soft (polymer content)  
1.20 % of 2-amino-2-methyl-1-propanol  
0.20 % of Cremophor A 25  
0.10 % of Uvinul P 25 PEG-25 PABA  
q.s. preservative  
10 q.s. perfume oil  
ad 100 % with water

Example 40:

Pump spray:

- 15 4.00 % of copolymer from Example 17  
1.00 % of panthenol  
0.10 % of Uvinul MS 40 Benzophenone-4  
q.s. preservative  
20 q.s. perfume oil  
ad 100 % with water

Example 41:

Pump spray:

- 25 4.00 % of copolymer from Example 9  
1.00 % of panthenol  
0.10 % of Uvinul M 40 Benzophenone-3  
q.s. preservative  
30 q.s. perfume oil  
ad 100 % with ethanol

Example 42:

Hair spray:

- 35 5.00 % of copolymer from Example 6  
0.10 % of silicone oil Dow Corning DC 190 Dimethicone Copolyol  
35.00 % of dimethyl ether  
5.00 % of n-pentane  
40 ad 100 % with ethanol  
q.s. perfume oil



Example 43:

Hairspray VOC 55 %:

3.00 % of copolymer from Example 4

5 7.00 % of Luviset P.U.R.

Polyurethane-1

40.00 % of dimethyl ether

15.00 % of ethanol

q.s. perfume oil

ad 100 % with water

10

Example 44:

Hair gel:

0.5 % of Carbopol 980

Carbomer

15 3.00 % of copolymer from example 18

0.10 % of phytantriol

0.50 % of panthenol

q.s. of perfume oil

q.s. of preservative

20 ad 100 % with water

Example 45:

Hair shampoo or shower gel

25 0.5 % of copolymer from example 33

40.00 % of Texapon NSO

Sodium Laureth Sulfate

5.00 % of Tego betaine L 7

Cocamidopropyl Betaine

5.00 % of Plantacare 2000

Decyl Glucoside

1.00 % of propylene glycol

30 q.s. of citric acid

q.s. of preservative

1.00 % of sodium chloride

ad 100 % with water

35

40

45


We claim:

1. The use of polymers obtainable by free-radical polymerization  
5 of

- a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in  
the presence of  
b) polyether-containing compounds and  
10 c) optionally one or more other copolymerizable monomers

and subsequent at least partial hydrolysis of the ester  
functions of the original monomers a), in hair cosmetic  
formulations.

- 15 2. The use of polymers as claimed in claim 1, wherein the  
polymers are obtainable by free-radical polymerization of  
a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in  
20 the presence of  
b) polyether-containing compounds of the formula I



30 in which the variables independently of one another have  
the following meanings:

R<sup>1</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-,  
polyalcohol radical;

35 R<sup>5</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

R<sup>2</sup> to R<sup>4</sup> are  
-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -CH<sub>2</sub>-CH(R<sup>6</sup>)-,  
-CH<sub>2</sub>-CHOR<sup>7</sup>-CH<sub>2</sub>-;

40 R<sup>6</sup> is C<sub>1</sub>-C<sub>24</sub>-alkyl;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>24</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

45 A is -C(=O)-O-, -C(=O)-B-C(=O)-O-,  
-C(=O)-NH-B-NH-C(=O)-O-;

B is  $-(CH_2)_t-$ , arylene, optionally substituted;

n is from 1 to 1000;

5 s is from 0 to 1000;

t is from 1 to 12;

u is from 1 to 5000;

10

v is from 0 to 5000;

w is from 0 to 5000;

15

x is from 0 to 5000;

y is from 0 to 5000;

z is from 0 to 5000;

20

and

c) optionally one or more other copolymerizable monomers

25

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

3. The use of polymers as claimed in claim 2, wherein the polymers are obtainable by free-radical polymerization of

30

a) at least one vinyl ester of  $C_1$ - $C_{24}$ -carboxylic acids in the presence of

35

b) polyether-containing compounds of the formula I having an average molecular weight of from 300 to 100000 (number average), in which the variables independently of one another have the following meanings:

40

$R^1$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-, polyalcohol radical;

$R^5$  is hydrogen,  $C_1$ - $C_{12}$ -alkyl,  $R^6$ -C(=O)-,  $R^6$ -NH-C(=O)-;

$R^2$  to  $R^4$  are

45

$-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH_2-CH(R^6)-$ ,  
 $-CH_2-CHOR^7-CH_2-$ ;

40

R<sup>6</sup> is C<sub>1</sub>-C<sub>12</sub>-alkyl;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>12</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

5 n is from 1 to 8;

s is 0;

u is from 2 to 2000;

10

v is from 0 to 2000;

w is from 0 to 2000;

15

and

c) optionally one or more other copolymerizable monomers

20

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

4. The use of polymers as claimed in claim 2, wherein the polymers are obtainable by free-radical polymerizable of

25

a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of

30

b) polyether-containing compounds of the formula I having an average molecular weight of from 500 to 50000 (number average), in which the variables independently of one another have the following meaning:

R<sup>1</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

35

R<sup>5</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

R<sup>2</sup> to R<sup>4</sup> are

-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -CH<sub>2</sub>-CH(R<sup>6</sup>)-,  
-CH<sub>2</sub>-CHOR<sup>7</sup>-CH<sub>2</sub>-;

40

R<sup>6</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, R<sup>6</sup>-C(=O)-, R<sup>6</sup>-NH-C(=O)-;

45

n is 1;

s is 0;

u is from 5 to 500;

5 v is from 0 to 500;

w is from 0 to 500;

and

10

c) optionally at least one or more other copolymerizable monomers

15

and subsequent at least partial hydrolysis of the ester functions of the original monomers a), in hair cosmetic formulations.

20

5. The use of polymers as claimed in claim 1, wherein the polymers are obtainable by free-radical polymerization of

a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of

25

b) polyether-containing silicone derivatives

and

30

c) optionally one or more other copolymerizable monomers

and subsequent at least partial hydrolysis of the ester function of the original monomers a).


35

6. The use of polymers as claimed in claim 5, wherein the polymers are obtainable by free-radical polymerization of

a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of

40





b) polyether-containing silicone derivatives of the formula II



45


where:

42



$R^{13}$  is a  $C_1$ - $C_{40}$  organic radical which can contain amino, carboxyl or sulfonate groups, or where  $e = 0$ , is also the anion of an inorganic acid,

and where the radicals  $R^8$  can be identical or different, and come either from the group of aliphatic hydrocarbons having from 1 to 20 carbon atoms, are cyclic aliphatic hydrocarbons having from 3 to 20 carbon atoms, are of an aromatic nature or are identical to  $R^{12}$ , where:



with the proviso that at least one of the radicals  $R^8$ ,  $R^9$  or  $R^{10}$  is a polyalkylene oxide-containing radical as defined above,


and  $f$  is an integer from 1 to 6,  
 $a$  and  $b$  are integers such that the molecular weight of the polysiloxane block is between 300 and 30000,  
 $c$  and  $d$  can be integers between 0 and 50, with the proviso that the sum  $c + d$  is greater than 0, and  $e$  is 0 or 1,

and

## 43

optionally one or more other copolymerizable monomers and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

- 5 7. The use of polymers as claimed in claim 6, wherein formula II has the following meaning:



- 15 8. The use as claimed in claim 1, wherein the polymers are obtainable by free-radical polymerization of
- 20 a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of
- b) polyether-containing compounds obtainable by reaction of polyethyleneimines with alkylene oxides
- 25 and
- c) optionally one or more other copolymerizable monomers
- and subsequent at least partial hydrolysis of the ester functions of the original monomers a).
- 30 9. The use of polymers as claimed in claim 8, wherein the alkylene oxides used are ethylene oxide, propylene oxide, butylene oxide and mixtures thereof.
- 35 10. The use of polymers as claimed in claims 8 and 9, wherein the alkylene oxide used is ethylene oxide.
- 40 11. The use of polymers as claimed in claims 8, 9 and 10, wherein the polyethyleneimine has a molecular weight between 300 and 20000.
- 45 12. The use of polymers as claimed in claim 1, wherein the polyether-containing compounds b) have been prepared by polymerization of ethylenically unsaturated alkylene oxide-containing monomers and optionally other copolymerizable monomers.

13. The use of polymers as claimed in claim 12, wherein the polyether-containing compounds b) have been prepared by polymerization of polyalkylene oxide vinyl ethers and optionally other copolymerizable monomers.

5

14. The use of polymers as claimed in claim 12, wherein the polyether-containing compounds b) have been prepared by polymerization of polyalkylene oxide (meth)acrylates and optionally other copolymerizable monomers.

10

15. The use of polymers as claimed in claims 1 to 14, wherein c) is chosen from the group:

- 15 acrylic acid, methacrylic acid, maleic acid, fumaric acid, crotonic acid, maleic anhydride and its half-esters, methyl acrylate, methyl methacrylate, ethyl acrylate, ethyl methacrylate, n-butyl acrylate, n-butyl methacrylate, t-butyl acrylate, t-butyl methacrylate, isobutyl acrylate, isobutyl methacrylate, 2-ethylhexyl acrylate, stearyl acrylate, 20 stearyl methacrylate, N-t-butylacrylamide, N-octylacrylamide, 2-hydroxyethyl acrylate, hydroxypropyl acrylates, 2-hydroxyethyl methacrylate, hydroxypropyl methacrylates, alkylene glycol (meth)acrylates, styrene, unsaturated sulfonic acids such as, for example, acrylamidopropane 25 sulfonic acid, vinyl pyrrolidone, vinyl caprolactam, vinyl ethers, (e.g. methyl, ethyl, butyl or dodecyl vinyl ethers), vinylformamide, vinylmethacrylamide, vinylamine, 1-vinylimidazole, 1-vinyl-2-methylimidazole, N,N-dimethylaminomethyl methacrylate and 30 N-[3-(dimethylamino)propyl]methacrylamide; 3-methyl-1-vinylimidazolium chloride, 3-methyl-1-vinylimidazolium methylsulfate, N,N-dimethylaminoethyl methacrylate, N-[3-(dimethylamino)propyl]methacrylamide quaternized with 35 methyl chloride, methyl sulfate or diethyl sulfate.

16. The use of polymers as claimed in claims 1 and 15, wherein the quantitative ratios are

- 40 a) 10 - 90 % by weight  
b) 2 - 90 % by weight  
c) 0 - 50 % by weight.

17. The use of polymers as claimed in claims 1 to 15, wherein the 45 quantitative ratios are

- a) 50 - 97 % by weight



- b) 3 - 50 % by weight
- c) 0 - 30 % by weight.

18. The use of polymers as claimed in claims 1 to 15, wherein the  
5 quantitative ratios are

- a) 60 - 97 % by weight
- b) 3 - 40 % by weight
- c) 0 - 20 % by weight.

10

19. The use as claimed in claims 1 to 18, where a crosslinking is  
carried out after the hydrolysis.

20. The use as claimed in claim 19, where the crosslinking is  
15 carried out by aldehydes, dialdehydes or borates.

21. A hair cosmetic formulation which has the following  
composition:

- 20 a) 0.05 - 20 % by weight of the polymer as in claim 1
- b) 20 - 99.95 % by weight of water and/or alcohol
- c) 0 - 79.05 % by weight of other constituents.

22. A hair cosmetic formulation which has the following  
25 composition:

- a) 0.1 - 10 % by weight of the polymer as in claim 1
- b) 20 - 99.9 % by weight of water and/or alcohol
- c) 0 - 70 % by weight of a propellant
- 30 d) 0 - 20 % by weight of other constituents.

23. A hair cosmetic formulation which has the following  
composition:

- 35 a) 0.1 - 10 % by weight of the polymer as in claim 1
- b) 55 - 94.8 % by weight of water and/or alcohol
- c) 5 - 20 % by weight of a propellant
- d) 0.1 - 5 % by weight of an emulsifier
- e) 0 - 10 % by weight of other constituents.

40

24. A hair cosmetic formulation which has the following  
composition:

- a) 0.1 - 10 % by weight of the polymer as in claim 1
- 45 b) 60 - 99.85 % by weight of water and/or alcohol
- c) 0.05 - 10 % by weight of a gel former

d) 0 - 20 % by weight of other constituents.

25. A hair cosmetic formulation which has the following composition:

5

- a) 0.05 - 10 % by weight of the polymer as in claim 1,
- b) 25 - 94.95 % by weight of water
- c) 5 - 50 % by weight of surfactants
- d) 0 - 5 % by weight of another conditioning agent
- 10 e) 0 - 10 % by weight of other cosmetic constituents.

26. A polymer obtainable by free-radical polymerization of

15

- a) at least one vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid, in the presence of
- b) polyether-containing silicone derivatives and
- c) optionally one or more other copolymerizable monomers

20

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

27. A polymer obtainable by free-radical polymerization of

25

- a) a vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid in the presence of
- b) polyether-containing compounds obtainable by reaction of polyethyleneimines with alkylene oxides and
- c) optionally one or more other copolymerizable monomers

30

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

28. A polymer obtainable by free-radical polymerization of

35

- a) a vinyl ester of a C<sub>1</sub>-C<sub>24</sub> carboxylic acid in the presence of
- b) homo- and copolymers of ethylenically unsaturated polyether-containing compounds and
- c) optionally one or more other copolymerizable monomers

40

and subsequent at least partial hydrolysis of the ester functions of the original monomers a).

45

29. A crosslinked polymer obtainable by free-radical polymerization of

- 5           a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub> carboxylic acids in  
          the presence of  
          b) polyether-containing compounds and  
          c) optionally one or more further copolymerizable monomers

10           and subsequent at least partial hydrolysis of the ester  
          functions of the original monomers a), where the crosslinker  
          used is either already present during the polymerization, or  
          is added after the polymerization and hydrolysis.

15           30. A crosslinked polymer as claimed in claim 29, where the  
          crosslinkers used are aldehydes, dialdehydes or borates.

20           31. The crosslinked polymer as claimed in claim 29, wherein the  
          crosslinker is already present during the polymerization.

25

30

35

40

45

Hair cosmetic formulations

Abstract

5

The use of polymers obtainable by free-radical polymerization of

- 10
- a) at least one vinyl ester of  $C_1$ - $C_{24}$ -carboxylic acids in the presence of
  - b) polyether-containing compounds and
  - c) optionally one or more other copolymerizable monomers

15 and subsequent at least partial hydrolysis of the ester functions of the original monomers a), in hair cosmetic formulations.

20

25

30

35

40

45

**PCT**  
 WELTORGANISATION FÜR GEISTIGES EIGENTUM  
 Internationales Büro  
 INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE  
 INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)



<b>(51) Internationale Patentklassifikation <sup>7</sup> :</b> <p style="text-align: center; margin-top: 10px;"><b>A61K 7/06</b></p>	<b>A1</b>	<b>(11) Internationale Veröffentlichungsnummer:</b> <b>WO 00/49998</b>  <b>(43) Internationales Veröffentlichungsdatum:</b> 31. August 2000 (31.08.00)
<b>(21) Internationales Aktenzeichen:</b> PCT/EP00/01070 <b>(22) Internationales Anmeldedatum:</b> 10. Februar 2000 (10.02.00)  <b>(30) Prioritätsdaten:</b> 199 07 587.5      22. Februar 1999 (22.02.99)      DE  <b>(71) Anmelder (für alle Bestimmungsstaaten ausser US):</b> BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE).  <b>(72) Erfinder; und</b> <b>(75) Erfinder/Anmelder (nur für US):</b> DIEING, Reinhold [DE/DE]; Drosselstr. 5 A, D-67105 Schifferstadt (DE). GOTSCHKE, Michael [DE/DE]; Warmweiherstr. 15, D-52066 Aachen (DE). HÖSSEL, Peter [DE/DE]; Birkenweg 10, D-67105 Schifferstadt (DE). SANNER, Axel [DE/DE]; Lorsche Ring 2c, D-67227 Frankenthal (DE). LEINENBACH, Alfred [DE/DE]; Gartenweg 3, D-67161 Gönheim (DE).  <b>(74) Gemeinsamer Vertreter:</b> BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE).	<b>(81) Bestimmungsstaaten:</b> CA, JP, MX, US, europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Veröffentlicht</b> <i>Mit internationalem Recherchenbericht.</i>	
<b>(54) Title:</b> FORMULATIONS OF HAIR COSMETICS <b>(54) Bezeichnung:</b> HAARKOSMETISCHE FORMULIERUNGEN  <b>(57) Abstract</b> <p>The invention relates to the use of polymers which can be obtained by radical polymerisation of a) at least one vinyl ester of C<sub>1</sub>-C<sub>24</sub>-carboxylic acids in the presence of b) compounds containing polyether and c) optionally, one or more other copolymerisable monomers, followed by the at least partial saponification of the ester functions of the original monomers a), in formulations of hair cosmetics.</p> <b>(57) Zusammenfassung</b> <p>Verwendung von Polymerisaten, die erhältlich sind durch radikalische Polymerisation von a) mindestens einem Vinylester von C<sub>1</sub>-C<sub>24</sub>-Carbonsäuren in Gegenwart von b) polyetherhaltigen Verbindungen und c) gegebenenfalls eines oder mehreren weiteren copolymerisierbaren Monomeren und anschließender zumindest teilweiser Verseifung der Esterfunktionen der ursprünglichen Monomeren a), in haarkosmetischen Formulierungen.</p>		

# Declaration, Power of Attorney

Page 1 of 4

0050/049774

We (I), the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Hair cosmetic formulations

the specification of which

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as

Application Serial No. \_\_\_\_\_

and amended on \_\_\_\_\_.

☒ was filed as PCT international application

Number PCT/EP00/01070

on February 10, 2000

and was amended under PCT Article 19

on \_\_\_\_\_ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
19907587.5	Germany	22 February 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Codes, § 119(e) of any United States provisional application(s) listed below.

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.

Filing Date

Status (pending, patented,  
abandoned)

_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

2-

And we (I) hereby appoint **Messrs. HERBERT. B. KEIL**, Registration Number 18,967; and **RUSSEL E. WEINKAUF**, Registration Number 18,495; the address of both being Messrs. Keil & Weinkauff, 1101 Connecticut Ave., N.W., Washington, D.C. 20036 (telephone 202-659-0100), our attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to sign the drawings, to receive the patent, and to transact all business in the Patent Office connected therewith.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

100 Reinhold Dieing  
NAME OF INVENTOR

Reinhold Dieing  
Signature of Inventor

Date February 24, 2000

Drosselstr. 5 A  
67105 Schifferstadt  
Germany DEX  
Citizen of: Germany  
Post Office Address: same as residence

200 Michael Gotsche  
NAME OF INVENTOR

Michael Gotsche  
Signature of Inventor

Date February 24, 2000

Warmweiherstr. 15  
52066 Aachen  
Germany DEX  
Citizen of: Germany  
Post Office Address: same as residence

1012000 Peter Hössel  
NAME OF INVENTOR

Peter Hössel  
Signature of Inventor

Date February 24, 2000

Birkenweg 10  
67105 Schifferstadt  
Germany DEX  
Citizen of: Germany  
Post Office Address: same as residence

400 Axel Sanner  
NAME OF INVENTOR

Axel Sanner  
Signature of Inventor

Date February 24, 2000

Lorscher Ring 2c  
67227 Frankenthal  
Germany DEX  
Citizen of: Germany  
Post Office Address: same as residence



Declaration

Page 4 of 4

0050/049774

500 Alfred Leinenbach  
NAME OF INVENTOR

Alfred Leinenbach  
Signature of Inventor

Date February 24, 2000

Gartenweg 3  
67161 Gönheim  
Germany DEX  
Citizen of: Germany  
Post Office Address: same as residence

600 Keith Leslie Rutherford  
NAME OF INVENTOR

Keith Leslie Rutherford  
Signature of Inventor

Date August 2, 2001

21 Lower Road  
Wirral CH62 5EG  
Great Britain GB  
Citizen of: Great Britain  
Post Office Address: same as residence

TO THE OFFICE OF THE SECRETARY OF COMMERCE